

À des fins de recherche uniquement

Anticorps Monoclonal anti-CD107a / LAMP1



Numéro de catalogue: 65050-1-Ig 2 Publications

Informations de base

Numéro de catalogue:	BC006785	Méthode de purification:
65050-1-Ig	Purification par affinité	
Taille:	Identification du gène (NCBI):	CloneNo.:
500ug , 0.5 mg/ml	16783	1D4B
Hôte:	Nom complet:	Dilutions recommandées:
Rat	lysosomal-associated membrane protein 1	IF 1:50-1:500
Isotype:		
IgG2a		

Applications

Applications testées:	Contrôles positifs:
FC, IF	IF : cellules NIH/3T3,
Demandes citées:	
IF	
Spécificité de l'espèce:	
souris	
Espèces citées:	
souris	

Informations générales

LAMP1 (CD107a) is a heavily glycosylated membrane protein enriched in the lysosomal membrane. LAMP1 is extensively glycosylated with asparagine-linked oligosaccharides which protect it from intracellular proteolysis (PMID: 10521503). Although LAMP1 is expressed largely in the endosome-lysosomal membrane of cells, it is also found on the plasma membrane (PMID: 16168398). Elevated LAMP1 expression at the cell surface has also been detected during platelet and granulocytic cell activation, as well as in some tumor cells (PMID: 29085473). LAMP1 functions to provide selectins with carbohydrate ligands. This protein has also been shown to be a marker of degranulation on lymphocytes such as CD8+ and NK cells and may also play a role in tumor cell differentiation and metastasis (PMID: 18835598; 29085473; 9426697).

Publications notables

Autrice	Pubmed ID	Journal	Application
Weidong Le	37292937	Res Sq	IF
Rui Hua	36924084	J Cell Physiol	IF

Stockage

Stockage:
Store at 2-8°C. Stable for one year after shipment.


Tampon de stockage:
Solution saline tamponnée au borate, pH 8.2.

For technical support and original validation data for this product please contact:
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



1X10⁶ NIH/3T3 cells were intracellularly stained with 0.50 ug Anti-Mouse CD107a (65050-1-Ig, Clone: 1D4B) (red) or 0.50 ug isotype control antibody (blue), and Fluorescein (FITC)-conjugated Affinipure Goat Anti-Rat IgG(H+L) with dilution of 1:1000. Cells were fixed with 90% MeOH.

Immunofluorescent analysis of (4% PFA) fixed NIH/3T3 cells using 65050-1-Ig (CD107a antibody) at dilution of 1:100 and Rhodamine (TRITC)-conjugated Goat Anti-Rat IgG(H+L).