À des fins de recherche uniquement

Anticorps Monoclonal anti-CD16

Numéro de catalogue:65090-1-lg



Méthode de purification:

Purification par affinité

CloneNo.:

3G8

Informations de base

Informations générales

Numéro de catalogue:

65090-1-lg

Taille:

100ug , 0.5 mg/ml

Hôte:

Mouse

Isotype: IgG1, kappa

Spécificité de l'espèce:

Humain

Applications testées:

CD16 is a 50-70-kDa low affinity Fc receptor found on the surface of natural killer cells, neutrophil

 $polymorphonuclear\ leukocytes, monocytes\ and\ macrophages.\ CD16\ mediates\ antibody-dependent\ cellular$ $cytotoxicity \ (ADCC) \ and \ other \ antibody-dependent \ responses, such as \ phagocytosis. \ CD16 \ has \ been \ identified \ as \ Fc$ receptors Fcy RIIIa (CD16a) and Fcy RIIIb (CD16b), encoded by two nearly identical genes, FCGR3A and the FCGR3B. Clone 3G8 recognizes both the CD16a and CD16b (PMID: 7592758).

Numéro d'acquisition GenBank:

Identification du gène (NCBI):

Fc fragment of IgG, low affinity IIIa,

BC017865

Nom complet:

receptor (CD16a)

MW calculé 254 aa, 29 kDa

Stockage

Applications

Store at 2-8°C. Stable for one year after shipment.

Tampon de stockage:


PBS with 0.09% sodium azide.

in USA), or 1(312) 455-8498 (outside USA)


W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



1X10^6 human PBMCs were surface co-stained with FITC Anti-Human CD3 and 0.2 ug Anti-Human CD16 (65090-1-Ig, Clone:3G8) and APC-Goat anti-Mouse IgG at dilution 1:1000. Cells were not fixed. Lymphocytes were gated.



1X10^6 human PBMCs were surface stained with 0.2 ug Anti-Human CD16 (65090-1-lg, Clone:3G8) and APC-Goat anti-Mouse IgG at dilution 1:1000. Cells were not fixed. Lymphocytes were gated.