## À des fins de recherche uniquement

## Anticorps Monoclonal anti-CD3

Numéro de catalogue:65133-1-lg



Méthode de purification:

Purification par affinité

CloneNo.:

OKT3

Informations de base

Numéro de catalogue:

65133-1-lg

Taille:

100ug , 0.5 mg/ml

Hôte:

Mouse

Isotype:

IgG2a, kappa MW calculé 207 aa, 23 kDa

**Applications** 

Applications testées:

Spécificité de l'espèce:

Humain

Informations générales

CD3 is a multimeric protein associated with the T-cell receptor (TCR) to form a complex involved in antigen recognition and signal transduction (PMID: 15885124). CD3 is composed of CD3 $\gamma$ ,  $\delta$ ,  $\epsilon$ , and  $\zeta$  chains (PMID: 1826255). It is expressed by thymocytes in a developmentally regulated manner, T cells, and some NK cells (PMID: 3289580). The TCR recognizes antigens bound to major histocompatibility complex (MHC) molecules. TCR-mediated peptide-MHC recognition is transmitted to the CD3 complex, leading to the intracellular signal transduction (PMID: 11985657).

Numéro d'acquisition GenBank:

Identification du gène (NCBI):

CD3e molecule, epsilon (CD3-TCR

BC049847

Nom complet:

complex)

Stockage


Stockage:

Store at 2-8°C. Stable for one year after shipment.


Tampon de stockage

PBS with 0.09% sodium azide.

## Données de validation sélectionnées



1X10^6 human PBMCs were surface co-stained with FITC Anti-human CD19 and 0.2 ug Anti-Human CD3 (65133-1-lg, Clone: OKT3) labeled with FlexAble CoraLite® Plus 647 Antibody Labeling Kit for Mouse IgG2a (KFA043) or CoraLite® Plus 647 conjugated Mouse IgG2a Isotype Control. Cells were not fixed. Lymphocytes were gated.



1X10^6 human PBMCs were surface stained with 0.2 ug Anti-Human CD3 (65133-1-1g, Clone: OKT3) labeled with FlexAble CoraLite® Plus 647 Antibody Labeling Kit for Mouse IgCJa (KFAO43) (red) or CoraLite® Plus 647 conjugated Mouse IgCJa Isotype Control. Cells were not fixed. Lymphocytes were gated.