

À des fins de recherche uniquement

# Anticorps Monoclonal anti-RNH1

Numéro de catalogue: 66028-1-Ig **Phare**



## Informations de base

<b>Numéro de catalogue:</b> 66028-1-Ig	<b>Numéro d'acquisition GenBank:</b> BC000677	<b>Méthode de purification:</b> Purification par protéine A
<b>Taille:</b> 150ul , Concentration: 1180 µg/ml by Bradford method using BSA as the standard;	<b>Identification du gène (NCBI):</b> 6050	<b>CloneNo.:</b> 4A1C4
<b>Hôte:</b> Mouse	<b>Nom complet:</b> ribonuclease/angiogenin inhibitor 1	<b>Dilutions recommandées:</b> WB 1:1000-1:8000
<b>Isotype:</b> IgG2b	<b>MW calculé</b> 50 kDa	
<b>Immunogen Catalog Number:</b> AG18093	<b>MW observés:</b> 50 kDa	

## Applications

<b>Applications testées:</b> WB,ELISA	<b>Contrôles positifs:</b> WB : cellules HeLa, cellules Jurkat, cellules L02
<b>Spécificité de l'espèce:</b> Humain	

## Informations générales

Ribonuclease/ angiogenin inhibitor (RNH, synonyms: RAI, MGC4569, MGC18200, MGC54054) is a protein that binds tightly to ribonucleases in cells and may be essential in the control of mRNA degradation and gene expression. The human RNH gene has been regionally localized to chromosome band 11p15 by in situ hybridization.

## Stockage

**Stockage:**  
Stocker à -20°C. Stable pendant un an après l'expédition.  
**Tampon de stockage:**  
PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3  
L'aliquotage n'est pas nécessaire pour le stockage à -20C

**\*\*\* Les 20ul contiennent 0,1% de BSA.**


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**

## Données de validation sélectionnées



HeLa, Jurkat, L02 cells were subjected to SDS PAGE followed by western blot with 66028-1-Ig (RNH1 Antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.



WB result of RNH1 antibody (66028-1-Ig, 1:4000) with si-control and si-RNH1 transfected HeLa cells.