

À des fins de recherche uniquement

Anticorps Monoclonal anti-IL-29

Numéro de catalogue: 66163-1-Ig



Informations de base

Numéro de catalogue: 66163-1-Ig	Numéro d'acquisition GenBank: BC074985	Méthode de purification: Purification par protéine G
Taille: 150ul , Concentration: 1207 µg/ml by Bradford method using BSA as the standard;	Identification du gène (NCBI): 282618	CloneNo.: 1H3A3
Hôte: Mouse	Nom complet: interleukin 29 (interferon, lambda 1)	Dilutions recommandées: WB 1:5000-1:50000
Isotype: IgG1	MW calculé 200 aa, 22 kDa	
Immunogen Catalog Number: AG12444	MW observés: 22 kDa	

Applications

Applications testées:
WB, ELISA
Spécificité de l'espèce:
Humain

Contrôles positifs:
WB : Protéine recombinante, Protéine recombinante

Informations générales

Interleukin-29 (IL-29) is a cytokine belonging to the Type III interferon family, which has another two subfamilies IL-28A and IL-28B. They are also known as IFN- λ 1, IFN- λ 2 and IFN- λ 3, respectively. IL-29 is produced predominantly by maturing dendritic cells and macrophages. IL-29 plays an important role in the immune response against pathogens and especially against viruses by mechanisms similar to type I interferons. IL-29 receptor signals through JAK-STAT pathways leading to activated expression of interferon-stimulated genes and production of antiviral proteins.

Stockage

Stockage:
Stocker à -20°C. Stable pendant un an après l'expédition.
Tampon de stockage:
PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3
L'aliquotage n'est pas nécessaire pour le stockage à -20C

*** Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



GST tagged recombinant IL29 protein were subjected to SDS PAGE followed by western blot with 66163-1-Ig (IL-29 antibody at dilution of 1:20000 incubated at room temperature for 1.5 hours.