

À des fins de recherche uniquement

Anticorps Monoclonal anti-CD71

Numéro de catalogue: 66180-1-Ig

Phare

13 Publications



Informations de base

Numéro de catalogue:

66180-1-Ig

Numéro d'acquisition GenBank:

BC001188

Méthode de purification:

Purification par protéine A

Taille:

150ul, Concentration: 3400 µg/ml by Nanodrop and 1800 µg/ml by Bradford method using BSA as the standard;

Identification du gène (NCBI):

7037

CloneNo.:

3C11F11

Hôte:

Mouse

Nom complet:

transferrin receptor (p90, CD71)

Dilutions recommandées:

WB 1:5000-1:50000

IP 0.5-4.0 ug for IP and 1:500-1:2000 for WB

Isotype:

IgG2a

MW calculé

85 kDa

IHC 1:2000-1:8000

IF 1:200-1:800

Immunogen Catalog Number:

AG21612

Applications

Applications testées:

FC, IF, IHC, IP, WB, ELISA

Demandes citées:

Blocking assay, IF, WB

Spécificité de l'espèce:

Humain

Espèces citées:

Humain, souris

Remarque-IHC: il est suggéré de démasquer l'antigène avec un tampon de TE buffer pH 9,0; (*) A défaut, 'le démasquage de l'antigène peut être 'effectué avec un tampon citrate pH 6,0.

Contrôles positifs:

WB : cellules U-251, cellules A431, cellules A549, cellules HeLa, cellules HepG2, cellules K-562, cellules Raji, cellules TF-1, cellules U2OS

IP : cellules HeLa,

IHC : tissu placentaire humain, tissu de cancer du sein humain

IF : tissu de cancer du sein humain,

Informations générales

CD71, also known as transferrin receptor protein 1 (TfR1), is a transmembrane glycoprotein composed of two disulfide-linked monomers, each of 90 kDa molecular weight. Each monomer binds one holo-transferrin molecule creating an iron-Tf-TfR complex which enters the cell by endocytosis. CD71 is present on actively proliferating cells and is essential for iron transport into proliferating cells.

Publications notables

Autrice	Pubmed ID	Journal	Application
Ping Han	36309297	Free Radic Biol Med	WB
Lingjuan Wang	34732689	Cell Death Discov	WB
Yangli Tian	34081703	PLoS Genet	WB

Stockage

Stockage:

Stocker à -20°C. Stable pendant un an après l'expédition.

Tampon de stockage:

PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3

L'aliquotage n'est pas nécessaire pour le stockage à -20C

*** Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Données de validation sélectionnées




Untreated and PNGase F-treated lysates of Raji cells and HeLa cells were subjected to SDS PAGE followed by western blot with 66180-1-Ig (CD71 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. PNGase F was obtained from Atagenix (cat.NO. ata808).




Various lysates were subjected to SDS PAGE followed by western blot with 66180-1-Ig (CD71 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Beta Actin Monoclonal antibody (HRP-66009) as loading control.




1X10⁶ Raji cells were stained with 0.2ug CD71 antibody (66180-1-Ig, red) and control antibody (blue). Fixed with 4% PFA blocked with 3% BSA (30 min). FITC-Goat anti-mouse IgG with dilution 1:100.




Immunohistochemical analysis of paraffin-embedded human placenta tissue slide using 66180-1-Ig (CD71 antibody) at dilution of 1:4000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human placenta tissue slide using 66180-1-Ig (CD71 antibody) at dilution of 1:4000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed human breast cancer tissue using CD71 antibody (66180-1-Ig, Clone: 3C11F11) at dilution of 1:400 and CoraLite@488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



IP Result of anti-CD71 (IP:66180-1-Ig, 5ug; Detection:66180-1-Ig 1:1000) with HeLa cells lysate 2400ug.