

À des fins de recherche uniquement

Anticorps Monoclonal anti-CD43

Numéro de catalogue: 66224-1-Ig



Informations de base

Numéro de catalogue: 66224-1-Ig	Numéro d'acquisition GenBank: BC012350	Méthode de purification: Purification par protéine A
Taille: 150ul, Concentration: 3300 µg/ml by Nanodrop and 2000 µg/ml by Bradford method using BSA as the standard;	Identification du gène (NCBI): 6693	CloneNo.: 2A11D6
Hôte: Mouse	Nom complet: sialophorin	Dilutions recommandées: WB 1:500-1:2000 IHC 1:4000-1:16000 IF 1:200-1:800
Isotype: IgG2b	MW calculé: 400 aa, 43 kDa	
Immunogen Catalog Number: AG23595	MW observés: 115 kDa	

Applications

Applications testées:
FC, IF, IHC, WB, ELISA

Spécificité de l'espèce:
Humain

Remarque-IHC: il est suggéré de démasquer l'antigène avec un tampon de TE buffer pH 9.0; (*) À défaut, le démasquage de l'antigène peut être effectué avec un tampon citrate pH 6,0.

Contrôles positifs:

WB : cellules Jurkat, cellules K-562

IHC : tissu d'amygdalite humain, tissu d'appendicite humain, tissu de côlon humain

IF : cellules Jurkat, tissu d'amygdalite humain

Informations générales

CD43, also named leukosialin, sialophorin, galactoglycoprotein, leukocyte sialoglycoprotein, SPN, and is a mucin-like type I transmembrane protein. In humans, it is expressed in haematopoietic cells, including T lymphocytes, monocytes, granulocytes, natural killer cells, platelets, and haematopoietic stem cells, with the exception of mature erythrocytes and B cell subpopulations. The unglycosylated form of CD43 migrates on SDS-PAGE with a molecular weight of 54 kDa (PMID: 2943740), whereas CD43 from different haematopoietic cell lines displays increased molecular weights since it is glycosylated with O-linked chains that differ in core structure and sialylation.

Stockage

Stockage:

Stocker à -20°C. Stable pendant un an après l'expédition.

Tampon de stockage:

PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3

L'aliquotage n'est pas nécessaire pour le stockage à -20C

*** Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Données de validation sélectionnées




Jurkat cells were subjected to SDS PAGE followed by western blot with 66224-1-Ig (CD43 antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 66224-1-Ig (CD43 antibody) at dilution of 1:8000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 66224-1-Ig (CD43 antibody) at dilution of 1:8000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed Jurkat cells using 66224-1-Ig (CD43 antibody) at dilution of 1:400 and CoralLite488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



1X10⁶ Jurkat cells were intracellularly stained with 0.4 ug Anti-Human CD43 (66224-1-Ig, Clone:2A11D6) and CoralLite[®]488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Mouse IgG2b Isotype Control (66360-3-Ig, Clone: K11B8C4B5) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).