

À des fins de recherche uniquement

Anticorps Monoclonal anti-p38 MAPK



Numéro de catalogue: 66234-1-Ig

Phare

47 Publications

Informations de base

Numéro de catalogue: 66234-1-Ig	Numéro d'acquisition GenBank: BC031574	Méthode de purification: Purification par protéine A
Taille: 150ul, Concentration: 2400 µg/ml by Nanodrop and 1333 µg/ml by Bradford method using BSA as the standard;	Identification du gène (NCBI): 1432	CloneNo.: 1A1C2
Hôte: Mouse	Nom complet: mitogen-activated protein kinase 14	Dilutions recommandées: WB 1:2000-1:6000 IHC 1:250-1:1000
Isotype: IgG2b	MW calculé: 360 aa, 41 kDa	
Immunogen Catalog Number: AG5797	MW observés: 38-42 kDa	

Applications

Applications testées:

IHC, WB, ELISA

Demandes citées:

IHC, WB

Spécificité de l'espèce:

Humain, porc

Espèces citées:

Humain, porc, rat, souris

Remarque-IHC: il est suggéré de démasquer l'antigène avec un tampon de TE buffer pH 9,0; (*) A défaut, 'le démasquage de l'antigène peut être 'effectué avec un tampon citrate pH 6,0.

Contrôles positifs:

WB : cellules HeLa, cellules HEK-293, cellules HSC-T6, cellules Jurkat, cellules K-562, cellules MCF-7, cellules RAW 264.7, tissu cardiaque de porc, tissu cardiaque humain

IHC : tissu de cancer du poumon humain, tissu de gliome humain

Informations générales

MAPK14(mitogen-activated protein kinase 14) is also named as SAPIK2A, p38MAPK, CSBP1, RK, p38, EXIP, Mxi2, CSBP2, PRKM14, PRKM15, CSPB1, p38ALPHA and belongs to the MAP kinase subfamily. MAPK14-signaling is a central pathway for the integration of instructive signals in dendritic cells for T(H)17 differentiation and inflammation(PMID:22231518). It plays an important role in the regulation of hematopoietic stem cell self-renewal in vitro and inhibition of MAPK14 activation with a small molecule inhibitor may represent a novel approach to promote ex vivo expansion of hematopoietic stem cell(PMID:21198398). This protein has 4 isoforms produced by alternative splicing.

Publications notables

Autrice	Pubmed ID	Journal	Application
Ting Tang	33173989	Mol Med Rep	WB,IHC
Weiche Wu	30273672	Free Radic Biol Med	WB
Hongfei Zhou	36115171	Phytomedicine	WB

Stockage

Stockage:

Stocker à -20°C. Stable pendant un an après l'expédition.

Tampon de stockage:

PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3

L'aliquotage n'est pas nécessaire pour le stockage à -20C

*** Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Données de validation sélectionnées




human heart tissue were subjected to SDS PAGE followed by western blot with 66234-1-Ig (p38 MAPK Antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.




WB result of p38 MAPK antibody (66234-1-Ig; 1:1000; incubated at room temperature for 1.5 hours) with sh-Control and sh-p38 MAPK transfected HEK-293 cells.




Various lysates were subjected to SDS PAGE followed by western blot with 66234-1-Ig (p38 MAPK antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human lung cancer tissue slide using 66234-1-Ig (p38 MAPK antibody) at dilution of 1:500 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human lung cancer tissue slide using 66234-1-Ig (p38 MAPK antibody) at dilution of 1:500 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Various lysates were subjected to SDS PAGE followed by western blot with 66234-1-Ig (p38 MAPK antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.