

À des fins de recherche uniquement

Anticorps Monoclonal anti-OTUD7B



Numéro de catalogue: 66276-1-Ig

Phare

6 Publications

Informations de base

Numéro de catalogue: 66276-1-Ig	Numéro d'acquisition GenBank: BC020622	Méthode de purification: Purification par protéine A
Taille: 150ul, Concentration: 1300 µg/ml by Nanodrop and 1000 µg/ml by Bradford method using BSA as the standard;	Identification du gène (NCBI): 56957	CloneNo.: 2C12D9
Hôte: Mouse	Nom complet: OTU domain containing 7B	Dilutions recommandées: WB 1:500-1:2000
Isotype: IgG2b	MW calculé: 427aa,48 kDa; 843aa,93 kDa	
Immunogen Catalog Number: AG10513	MW observés: 92 kDa	

Applications

Applications testées: WB, ELISA	Contrôles positifs: WB: cellules Jurkat,
Demandes citées: IF, IHC, WB	
Spécificité de l'espèce: Humain	
Espèces citées: Humain	

Informations générales

OUTD7B is a member of the A20 family of the deubiquitinating cysteine proteases, which has deubiquitinating activity that is directed towards 'Lys-48' or 'Lys-63'-linked polyubiquitin chains. It plays a key role in the negative feedback regulation of NF- κ B signaling in response to multiple stimuli and has been described as central gatekeeper in inflammation and immunity.

Publications notables

Autrice	Pubmed ID	Journal	Application
Boxiang Zhang	33198776	J Exp Clin Cancer Res	IHC
Shuo Tian	32453962	Autophagy	WB, IF
Jianing Tang	34035221	Cell Death Dis	WB, IHC

Stockage


Stockage:
Stocker à -20°C. Stable pendant un an après l'expédition.
Tampon de stockage:
PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3
L'aliquotage n'est pas nécessaire pour le stockage à -20C

*** Les 20ul contiennent 0,1% de BSA.

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



Jurkat cells were subjected to SDS PAGE followed by western blot with 66276-1-Ig (OTUD7B Antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.