

À des fins de recherche uniquement

# Anticorps Monoclonal anti-PEX5

Numéro de catalogue: 66309-1-Ig

Phare

1 Publications



## Informations de base

Numéro de catalogue: 66309-1-Ig	Numéro d'acquisition GenBank: BC010621	Méthode de purification: Purification par protéine G
Taille: 150ul, Concentration: 1500 µg/ml by Nanodrop and 1000 µg/ml by Bradford method using BSA as the standard;	Identification du gène (NCBI): 5830	CloneNo.: 1C2C3
Hôte: Mouse	Nom complet: peroxisomal biogenesis factor 5	Dilutions recommandées: WB 1:500-1:2000 IHC 1:50-1:500
Isotype: IgG1	MW calculé: 631 aa, 70 kDa	
Immunogen Catalog Number: AG3268	MW observés: 70-80 kDa	

## Applications

### Applications testées:

IHC, WB, ELISA

### Demandes citées:

WB

### Spécificité de l'espèce:

Humain, porc, rat

### Espèces citées:

Humain

**Remarque-IHC: il est suggéré de démasquer l'antigène avec un tampon de TE buffer pH 9,0; (\*) A défaut, 'le démasquage de l'antigène peut être effectué avec un tampon citrate pH 6,0.**

### Contrôles positifs:

WB : tissu cérébral de porc, cellules HEK-293, tissu cérébral humain foetal

IHC : tissu hépatique humain, tissu rénal humain

## Informations générales

The peroxisomal targeting signal type1 (PTS1) receptor, PEX5, is one member of peroxins (PEXs) which are proteins required for peroxisome assembly. PEX5 and PEX7 function as receptors that recognize PTS1- and PTS2- containing proteins, respectively, and PEX5 binds PTS1 through its C-terminal 40-kDa tetratricopeptide repeat domain. It is a predominantly cytoplasmic, partly peroxisomal protein that appears to shuttle between these compartments as it mediates the import of PTS1-containing proteins. PEX5 has been reported to interact with PEX10, PEX12, PEX13, and PEX14. Defects in PEX5 are a cause of Zellweger syndrome (ZWS), which is a lethal peroxisome biogenesis disorder.

## Publications notables

Autrice	Pubmed ID	Journal	Application
Léa P Wilhelm	36215693	EMBO J	WB

## Stockage

### Stockage:

Stocker à -20°C. Stable pendant un an après l'expédition.

### Tampon de stockage:

PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3

L'aliquotage n'est pas nécessaire pour le stockage à -20C

\*\*\* Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Données de validation sélectionnées




WB result of PEX5 antibody (66309-1-Ig: 1:1000; incubated at room temperature for 1.5 hours) with sh-Control and sh-PEX5 transfected HEK-293 cells.



pig brain tissue were subjected to SDS PAGE followed by western blot with 66309-1-Ig (PEX5 Antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human liver tissue slide using 66309-1-Ig (PEX5 Antibody) at dilution of 1:200 (under 10x lens).



Immunohistochemical analysis of paraffin-embedded human liver tissue slide using 66309-1-Ig (PEX5 Antibody) at dilution of 1:200 (under 40x lens).