

À des fins de recherche uniquement

Anticorps Monoclonal anti-RSPO3

Numéro de catalogue: 66314-1-Ig



Informations de base

Numéro de catalogue:	BC022367	Méthode de purification:
66314-1-Ig		Purification par protéine A
Taille:	Identification du gène (NCBI):	CloneNo.:
150ul , Concentration: 1900 µg/ml by Nanodrop and 1000 µg/ml by Bradford method using BSA as the standard;	84870	1E4F1
Hôte:	Nom complet:	Dilutions recommandées:
Mouse	R-spondin 3 homolog (Xenopus laevis)	WB 1:1000-1:4000
Isotype:	MW calculé	IHC 1:50-1:500
IgG2b	272 aa, 31 kDa	IF 1:50-1:500
Immunogen Catalog Number:	MW observés:	
AG9899	32 kDa	

Applications

Applications testées:	Contrôles positifs:
IF, IHC, WB, ELISA	WB : tissu de muscle squelettique humain, cellules A549
Spécificité de l'espèce:	IHC : tissu de cancer du poumon humain,
Humain	IF : cellules A549,
Remarque-IHC: il est suggéré de démasquer l'antigène avec un tampon de TE buffer pH 9,0; (*) À défaut, le démasquage de l'antigène peut être effectué avec un tampon citrate pH 6,0.	

Informations générales

RSPO3, also named as R-spondin-3, is a 272 amino acid protein, which contains 1 TSP type-1 domain and 2 FU (furin-like) repeats. RSPO3 may be a secreted protein and belongs to the R-spondin family. RSPO3 is expressed at higher level in placenta, small intestine, fetal thymus and lymph node. RSPO3 is an activator of the canonical Wnt signaling pathway by acting as a ligand for LGR4-6 receptors. The four R-spondins (RSPO1-4) and three related leucinerich repeat-containing G-protein coupled receptors, LGR4, LGR5, and LGR6 (LGR4-6), constitute a ligand-receptor system that plays critical roles in development, stem cell survival, and oncogenesis. Ectopic expression of RSPO2/3 in mouse mammary epithelial cells led to an increase in invasiveness in vitro and in tumor formation and metastasis in vivo.

Stockage




Stockage:
Stocker à -20°C. Stable pendant un an après l'expédition.
Tampon de stockage:
PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3
L'aliquotage n'est pas nécessaire pour le stockage à -20C

*** Les 20ul contiennent 0,1% de BSA.

For technical support and original validation data for this product please contact:
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Données de validation sélectionnées



human skeletal muscle tissue were subjected to SDS PAGE followed by western blot with 66314-1-Ig (RSPO3 Antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.

Immunohistochemical analysis of paraffin-embedded human lung cancer tissue slide using 66314-1-Ig (RSPO3 Antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).

Immunohistochemical analysis of paraffin-embedded human lung cancer tissue slide using 66314-1-Ig (RSPO3 Antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed A549 cells using 66314-1-Ig(RSPO3 antibody) at dilution of 1:100 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Mouse IgG(H+L).