

À des fins de recherche uniquement

Anticorps Monoclonal anti-NONO

Numéro de catalogue: 66361-1-Ig **1 Publications**



Informations de base

Numéro de catalogue: 66361-1-Ig	Numéro d'acquisition GenBank: BC010049	Méthode de purification: Purification par protéine G
Taille: 150ul , Concentration: 1000 µg/ml by Bradford method using BSA as the standard;	Identification du gène (NCBI): 4841	CloneNo.: 2A2B10
Hôte: Mouse	Nom complet: non-POU domain containing, octamer-binding	Dilutions recommandées: WB 1:500-1:2000
Isotype: IgG1	MW calculé 54 kDa	
Immunogen Catalog Number: AG17609	MW observés: 54 kDa	

Applications

Applications testées: WB, ELISA	Contrôles positifs: WB : cellule C2C12, cellules ROS1728, tissu de muscle squelettique de souris
Demandes citées: WB	
Spécificité de l'espèce: Humain, rat, souris	
Espèces citées: souris	

Informations générales

Non-pou domain-containing octamer-binding protein(NONO), other named p54nrb, belongs to DBHS family, whose protein are predominately nuclear and involves in transcriptional and posttranscriptional gene regulatory functions as well as DNA repair. This family has roles in a wide range of biological processes, such as the regulation of circadian rhythm, carcinogenesis and progression of cancer. NONO forms heterodimer with SFPQ, and has been implicated in nuclear retention of A- to I-edited RNA as paraspeckle components, pre-mRAN 3'-end formation, cAMP cycling and transcriptional activation.

Publications notables

Autrice	Pubmed ID	Journal	Application
Nian Liu	35953472	Nat Commun	WB

Stockage


Stockage:
Stocker à -20°C. Stable pendant un an après l'expédition.
Tampon de stockage:
PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3
L'aliquotage n'est pas nécessaire pour le stockage à -20C

*** Les 20ul contiennent 0,1% de BSA.

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



C2C12 cell were subjected to SDS PAGE followed by western blot with 66361-1-Ig (NONO Antibody) at dilution of 1:1000 incubated at room temperature for 1.5 hours.