

À des fins de recherche uniquement

Anticorps Monoclonal anti-SIRT2

Numéro de catalogue: 66410-1-Ig

Phare

6 Publications



Informations de base

Numéro de catalogue: 66410-1-Ig	Numéro d'acquisition GenBank: BC003547	Méthode de purification: Purification par protéine G
Taille: 150ul, Concentration: 1400 µg/ml by Nanodrop and 1000 µg/ml by Bradford method using BSA as the standard;	Identification du gène (NCBI): 22933	CloneNo.: 1D8G10
Hôte: Mouse	Nom complet: sirtuin (silent mating type information regulation 2 homolog) 2 (S. cerevisiae)	Dilutions recommandées: WB 1:5000-1:50000 IHC 1:50-1:500 IF 1:50-1:500
Isotype: IgG1	MW calculé: 43 kDa	
Immunogen Catalog Number: AG7756	MW observés: 37-45 kDa	

Applications

Applications testées:

IF, IHC, WB, ELISA

Demandes citées:

IF, IHC, WB

Spécificité de l'espèce:

Humain, porc, rat, souris

Espèces citées:

Humain, rat, souris

Remarque-IHC: il est suggéré de démasquer l'antigène avec un tampon de TE buffer pH 9,0; (*) A défaut, 'le démasquage de l'antigène peut être 'effectué avec un tampon citrate pH 6,0.

Contrôles positifs:

WB : cellules LNCaP, cellules MCF-7, tissu cardiaque de porc, tissu cardiaque de rat, tissu cérébral de porc, tissu cérébral de rat, tissu cérébral de souris, tissu cérébral humain foetal

IHC : tissu rénal humain, tissu cardiaque humain

IF : cellules HepG2,

Informations générales

The silent information regulator(SIR2) family of genes are highly conserved from prokaryotes to eukaryotes and are involved in diverse processes, including transcriptional regulation, cell cycle progression, DNA-damage repair and aging. SIR2 contains a 323 amino acid catalytic core domain with a NAD-binding domain and a large groove which is the likely site of catalysis. SIR2 is widely expressed, highly expressed in heart, brain and skeletal muscle, while it is weakly expressed in placenta and lung. Down-regulated in many gliomas suggesting that it may act as a tumor suppressor gene in human gliomas possibly through the regulation of microtubule network. SIRT2 exists various isoforms and the molecular weight of isoforms are 35kDa, 40 kDa, and 42 kDa.

Publications notables

Autrice	Pubmed ID	Journal	Application
Rui Lin	36082450	Prostate	WB
Scarlett Acklin	35875690	Neurooncol Adv	IHC
David Siegel	33360352	Redox Biol	WB

Stockage

Stockage:

Stocker à -20°C. Stable pendant un an après l'expédition.

Tampon de stockage:

PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3

L'aliquoteage n'est pas nécessaire pour le stockage à -20C

*** Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Données de validation sélectionnées




Various lysates were subjected to SDS PAGE followed by western blot with 66410-1-Ig (SIRT2 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Alpha Tubulin Monoclonal antibody (HRP-66031) as loading control.



Immunohistochemical analysis of paraffin-embedded human kidney tissue slide using 66410-1-Ig (SIRT2 Antibody) at dilution of 1:200 (under 40x lens).



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using 66410-1-Ig (SIRT2 antibody) at dilution of 1:100 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



Various lysates were subjected to SDS PAGE followed by western blot with 66410-1-Ig (SIRT2 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.