

À des fins de recherche uniquement

Anticorps Monoclonal anti-PLEK

Numéro de catalogue: 66431-1-Ig **Phare**



Informations de base

Numéro de catalogue: 66431-1-Ig	Numéro d'acquisition GenBank: BC018549	Méthode de purification: Purification par protéine G
Taille: 150ul, Concentration: 1600 µg/ml by Nanodrop and 1000 µg/ml by Bradford method using BSA as the standard;	Identification du gène (NCBI): 5341	CloneNo.: 1D5C2
Hôte: Mouse	Nom complet: pleckstrin	Dilutions recommandées: WB 1:2000-1:20000 IHC 1:50-1:500 IF 1:200-1:800
Isotype: IgG1	MW calculé: 350 aa, 40 kDa	
Immunogen Catalog Number: AG3187	MW observés: 40-47 kDa	

Applications

Applications testées:

IF, IHC, WB, ELISA

Spécificité de l'espèce:

Humain, porc

Remarque-IHC: il est suggéré de démasquer l'antigène avec un tampon de TE buffer pH 9.0; (*) À défaut, le démasquage de l'antigène peut être effectué avec un tampon citrate pH 6,0.

Contrôles positifs:

WB : cellules HL-60, cellules THP-1, cellules U-937, tissu splénique de porc

IHC : tissu de cancer du côlon humain,

IF : tissu splénique de souris,

Informations générales

PLEK (pleckstrin; also known as p47) was originally identified as the major PKC substrate in platelets and later is found to be expressed in all cells of the hemopoietic system. Through two pleckstrin homology (PH) domains at its N and C termini, PLEK may interact with various protein and/or lipid ligands and serve as an intracellular adaptor/targeting protein. Predominantly cytosolic in unstimulated cells, PLEK would undergo transient redistribution to phagosomal membrane in response to stimuli. PLEK has been found to be hyperphosphorylated in diabetic mononuclear phagocytes and promote proinflammatory cytokine secretion in diabetes. (10477609, 17579087)

Stockage

Stockage:

Stocker à -20°C. Stable pendant un an après l'expédition.

Tampon de stockage:

PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3

L'aliquotage n'est pas nécessaire pour le stockage à -20C

*** Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Données de validation sélectionnées




WB result of PLEK antibody (66431-1-Ig; 1:2000; incubated at room temperature for 1.5 hours) with sh-Control and sh-PLEK transfected U-937 cells.




Immunofluorescent analysis of (4% PFA) fixed mouse spleen tissue using PLEK antibody (66431-1-Ig, Clone: 1D5C2) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).




Immunofluorescent analysis of (4% PFA) fixed mouse spleen tissue using PLEK antibody (66431-1-Ig, Clone: 1D5C2) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



HL-60 cells were subjected to SDS PAGE followed by western blot with 66431-1-Ig (PLEK Antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human colon cancer tissue slide using 66431-1-Ig (PLEK Antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human colon cancer tissue slide using 66431-1-Ig (PLEK Antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).