

À des fins de recherche uniquement

Anticorps Monoclonal anti-CCAR2

Numéro de catalogue: 66497-1-Ig **Phare**



Informations de base

Numéro de catalogue: 66497-1-Ig	Numéro d'acquisition GenBank: BC018269	Méthode de purification: Purification par protéine A
Taille: 150ul , Concentration: 1500 µg/ml by Nanodrop and 1000 µg/ml by Bradford method using BSA as the standard;	Identification du gène (NCBI): 57805	CloneNo.: 2D10F8
Hôte: Mouse	Nom complet: KIAA1967	Dilutions recommandées: WB 1:2000-1:16000 IHC 1:250-1:1000 IF 1:50-1:500
Isotype: IgG2a	MW calculé 923 aa, 103 kDa	
Immunogen Catalog Number: AG18374	MW observés: 130 kDa	

Applications

Applications testées:
FC, IF, IHC, WB, ELISA

Spécificité de l'espèce:
Humain, porc

Remarque-IHC: il est suggéré de démasquer l'antigène avec un tampon de TE buffer pH 9.0; (*) À défaut, le démasquage de l'antigène peut être effectué avec un tampon citrate pH 6.0.

Contrôles positifs:

WB : cellules HeLa, cellules HEK-293, cellules Jurkat, cellules MCF-7, tissu cérébral de porc

IHC : tissu de cancer du col de l'utérus humain, tissu de cancer du côlon humain, tissu de cancer du poumon humain, tissu de côlon humain

IF : cellules HeLa,

Informations générales

Cell cycle and apoptosis regulator protein 2 (CCAR2), previously known as DBC1/KIAA1967, regulates diverse cellular functions including transcription, mRNA splicing, metabolism, and circadian cycle. CCAR2 is a critical regulator of cell death or survival following cellular stresses. CCAR2 is a widely expressed protein involved in the regulation of a variety of transcriptional complexes. High expression of CCAR2 correlates with poor outcomes in many human tumor types such as squamous cell carcinoma (SCC).

Stockage

Stockage:
Stocker à -20°C. Stable pendant un an après l'expédition.
Tampon de stockage:
PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3
L'aliquotage n'est pas nécessaire pour le stockage à -20C

*** Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Données de validation sélectionnées




Various lysates were subjected to SDS PAGE followed by western blot with 66497-1-Ig (CCAR2 antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.




WB result of CCAR2 antibody (66497-1-Ig; 1:2000; incubated at room temperature for 1.5 hours) with sh-Control and sh-CCAR2 transfected HEK-293 cells.




Immunohistochemical analysis of paraffin-embedded human cervical cancer tissue slide using 66497-1-Ig (CCAR2 antibody) at dilution of 1:500 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human cervical cancer tissue slide using 66497-1-Ig (CCAR2 antibody) at dilution of 1:500 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HeLa cells using 66497-1-Ig (CCAR2 antibody) at dilution of 1:100 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



1×10^6 U2OS cells were intracellularly stained with 0.2 ug Anti-Human CCAR2 (66497-1-Ig, Clone:2D10F8) and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.2 ug Mouse IgG2a Isotype Control (66360-2-Ig, Clone: K11A1B2A2) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).