

À des fins de recherche uniquement

Anticorps Monoclonal anti-FOXC1

Numéro de catalogue: 66568-1-Ig **Phare**



Informations de base

Numéro de catalogue: 66568-1-Ig	Numéro d'acquisition GenBank: NM_001453	Méthode de purification: Purification par protéine G
Taille: 150ul, Concentration: 1500 µg/ml by Nanodrop and 1000 µg/ml by Bradford method using BSA as the standard;	Identification du gène (NCBI): 2296	CloneNo.: 1F4E11
Hôte: Mouse	Nom complet: forkhead box C1	Dilutions recommandées: WB 1:2000-1:20000
Isotype: IgG1	MW calculé 57 kDa	
	MW observés: 70-75 kDa	

Applications

Applications testées: WB, ELISA	Contrôles positifs:
Spécificité de l'espèce: Humain	WB : cellules HeLa, cellules HEK293, cellules HEK-293, cellules HepG2, cellules L02

Informations générales

FOXC1, also named as FKHL7 and FREAC3, binding of FREAC-3 and FREAC-4 to their cognate sites results in bending of the DNA at an angle of 80-90 degrees. Defects in FOXC1 are the cause of Axenfeld-Rieger syndrome type 3 (RIEG3). Defects in FOXC1 are the cause of iridogoniodysgenesis anomaly (IGDA). Defects in FOXC1 are a cause of Peters anomaly.

Stockage

Stockage:
Stocker à -20°C. Stable pendant un an après l'expédition.
Tampon de stockage:
PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3
L'aliquotage n'est pas nécessaire pour le stockage à -20C

***** Les 20ul contiennent 0,1% de BSA.**


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



Various cells were subjected to SDS PAGE followed by western blot with 66568-1-Ig (FOXC1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



WB result of FOXC1 antibody (66568-1-Ig; 1:10000; incubated at room temperature for 1.5 hours) with sh-Control and sh-FOXC1 transfected HEK-293 cells.