

À des fins de recherche uniquement

# Anticorps Monoclonal anti-CD100

Numéro de catalogue: 66582-1-Ig

Phare

1 Publications



## Informations de base

Numéro de catalogue:	66582-1-Ig	Numéro d'acquisition GenBank:	BC054500	Méthode de purification:	Purification par protéine A
Taille:	150ul , Concentration: 1300 µg/ml by 10507	Identification du gène (NCBI):	Nom complet:	CloneNo.:	1C1A6
	Nanodrop and 1000 µg/ml by Bradford method using BSA as the standard;	sema domain, immunoglobulin domain (Ig), transmembrane domain (TM) and short cytoplasmic domain, (semaphorin) 4D	Dilutions recommandées:	WB 1:1000-1:3000	IHC 1:225-1:900
Hôte:	Mouse	MW calculé			
Isotype:	IgG1	96 kDa	MW observés:	150 kDa	

## Applications

Applications testées:	Contrôles positifs:
IHC, WB, ELISA	WB : cellules HL-60, cellules HeLa, cellules K-562, cellules THP-1
Demandes citées:	IHC : tissu d'amygdale humain, tissu d'appendicite humain, tissu de lymphome humain, tissu rénal humain
WB	
Spécificité de l'espèce:	
Humain	
Espèces citées:	
Humain	
<b>Remarque-IHC: il est suggéré de démasquer l'antigène avec un tampon de TE buffer pH 9.0; (*) À défaut, le démasquage de l'antigène peut être effectué avec un tampon citrate pH 6,0.</b>	

## Informations générales

CD100 (also known as SEMA4D), a 150 kDa homodimer, belongs to the family of immune semaphorins, and CD100 is abundantly expressed on resting T cells, NK cells, and antigen-presenting cells. As a transmembrane glycoprotein, it is digested to its soluble form, sCD100, by specific matrix metalloproteinases. CD100 has been reported that the expression levels of CD100, and one of its receptors, plexin-B1 (PLXNB1), are increased in head and neck, prostate, colon, breast, and lung cancers, and the interaction between them provides oncogenic signaling essential for tumor growth and metastasis.

## Publications notables

Autrice	Pubmed ID	Journal	Application
Xiang Liu	36936030	Oncol Lett	WB

## Stockage

Stockage:  
Stocker à -20°C. Stable pendant un an après l'expédition.  
Tampon de stockage:  
PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3  
L'aliquotage n'est pas nécessaire pour le stockage à -20°C


\*\*\* Les 20ul contiennent 0,1% de BSA.

For technical support and original validation data for this product please contact:  
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free  
in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com  
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Données de validation sélectionnées



various cells were subjected to SDS PAGE followed by western blot with 66582-1-Ig (CD100 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 66582-1-Ig (CD100 antibody) at dilution of 1:450 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 66582-1-Ig (CD100 antibody) at dilution of 1:450 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).