

À des fins de recherche uniquement

Anticorps Monoclonal anti-c-Fos

Numéro de catalogue: 66590-1-Ig

Phare

52 Publications



Informations de base

Numéro de catalogue: 66590-1-Ig	Numéro d'acquisition GenBank: BC004490	Méthode de purification: Purification par protéine G
Taille: 150ul , Concentration: 1000 µg/ml by Nanodrop;	Identification du gène (NCBI): 2353	CloneNo.: 1G2C5
Hôte: Mouse	Nom complet: FOS	Dilutions recommandées: WB 1:5000-1:50000
Isotype: IgG1	MW calculé 41 kDa	
Immunogen Catalog Number: AG24340	MW observés: 55-60 kDa	

Applications

Applications testées:

FC, WB, ELISA

Demandes citées:

IHC, WB

Spécificité de l'espèce:

Humain, rat, souris

Espèces citées:

Humain, souris

Contrôles positifs:

WB : cellules HeLa, cellules HepG2, cellules Jurkat, cellules K-562, cellules RAW 264.7, cellules THP-1, cellules U-937

Informations générales

c-Fos, also named as FOS and G0/G1 switch regulatory protein 7, is a 380 amino acid protein, which contains 1 bZIP (basic-leucine zipper) domain and belongs to the bZIP family. c-Fos is expressed at very low levels in quiescent cells. When cells are stimulated to reenter growth, c-Fos undergo 2 waves of expression, the first one peaks 7.5 minutes following FBS induction. At this stage, the c-Fos protein is localized endoplasmic reticulum. The second wave of expression occurs at about 20 minutes after induction and peaks at 1 hour. At this stage, the c-FOS protein becomes nuclear. c-Fos is a very short-lived intracellular protein, which is very easy to degrade. The calculated molecular weight of c-Fos is 40 kDa, but Phosphorylated c-Fos protein is about 60-65 kDa. It is involved in important cellular events, including cell proliferation, differentiation and survival; genes associated with hypoxia; and angiogenesis; which makes its dysregulation an important factor for cancer development. It can also induce a loss of cell polarity and epithelial-mesenchymal transition, leading to invasive and metastatic growth in mammary epithelial cells. Expression of c-Fos is an indirect marker of neuronal activity because c-Fos is often expressed when neurons fire action potentials. Upregulation of c-Fos mRNA in a neuron indicates recent activity.

Publications notables

Autrice	Pubmed ID	Journal	Application
Ning Wang	36235607	Nutrients	WB
Xuming Wang	36187757	Front Physiol	WB,IHC
Hongbing Lin	36114617	Stem Cells Dev	WB

Stockage

Stockage:

Stocker à -20°C. Stable pendant un an après l'expédition.

Tampon de stockage:

PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3

L'aliquotage n'est pas nécessaire pour le stockage à -20C

*** Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Données de validation sélectionnées



WB result of c-Fos antibody (66590-1-Ig; 1:40000; incubated at room temperature for 1.5 hours) with sh-Control and sh-c-Fos transfected HepG2 cells.



Various lysates were subjected to SDS PAGE followed by western blot with 66590-1-Ig (c-Fos antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



1×10^6 HeLa cells were intracellularly stained with 0.5 ug Anti-Human c-Fos (66590-1-Ig, Clone:1G2C5) and CoraLite@488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.5 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).