

À des fins de recherche uniquement

Anticorps Monoclonal anti-NXN

Numéro de catalogue: 66603-1-Ig



Informations de base

Numéro de catalogue:	66603-1-Ig	Numéro d'acquisition GenBank:	BC009327	Méthode de purification:	Purification par protéine A
Taille:	150ul , Concentration: 1000 µg/ml by Bradford method using BSA as the standard;	Identification du gène (NCBI):	64359	CloneNo.:	2C6A4
Hôte:	Mouse	Nom complet:	nucleoredoxin	Dilutions recommandées:	WB 1:5000-1:50000 IHC 1:50-1:500 IF 1:400-1:1600
Isotype:	IgG1	MW calculé	435 aa, 48 kDa		
		MW observés:	48 kDa		
		Immunogen Catalog Number:	AG9248		

Applications

Applications testées:	IF, IHC, WB, ELISA	Contrôles positifs:	
Spécificité de l'espèce:	Humain, rat, souris	WB :	cellules HeLa, cellules HEK-293, cellules HepG2, cellules Jurkat
		IHC :	tissu rénal humain,
Remarque-IHC: il est suggéré de démasquer l'antigène avec un tampon de TE buffer pH 9,0; (*) À défaut, le démasquage de l'antigène peut être effectué avec un tampon citrate pH 6,0.			IF : cellules HEK-293,

Informations générales

Nucleoredoxin (NRV/NXN), a relatively uncharacterized member of the TRX family protein, has recently been reported to regulate the Wnt/β-catenin pathway, which itself regulates cell fate and early development, in a redox-dependent manner (PMID:17567240). Northern blot analysis detected several nucleoredoxin splice variants in adult mouse tissues, with highest expression in testis and skin and lowest expression in liver (PMID:9119370). It has 3 isoforms produced by alternative splicing. This antibody is specific to NXN.

Stockage

Stockage:
Stocker à -20°C. Stable pendant un an après l'expédition.
Tampon de stockage:
PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3
L'aliquotage n'est pas nécessaire pour le stockage à -20°C


*** Les 20ul contiennent 0,1% de BSA.

For technical support and original validation data for this product please contact:
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Données de validation sélectionnées




Immunofluorescent analysis of (-20°C Ethanol) fixed HEK-293 cells using NXN antibody (66603-1-Ig, Clone: 2C6A4) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



Immunohistochemical analysis of paraffin-embedded human kidney tissue slide using 66603-1-Ig (NXN antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human kidney tissue slide using 66603-1-Ig (NXN antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Various lysates were subjected to SDS PAGE followed by western blot with 66603-1-Ig (NXN antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control.