

À des fins de recherche uniquement

Anticorps Monoclonal anti-PLCB3

Numéro de catalogue: 66668-1-Ig

3 Publications



Informations de base

Numéro de catalogue:	66668-1-Ig	Numéro d'acquisition GenBank:	BC142681	Méthode de purification:	Purification par protéine G
Taille:	150ul , Concentration: 1100 µg/ml by 5331	Identification du gène (NCBI):	Nom complet: phospholipase C, beta 3 (phosphatidylinositol-specific)	CloneNo.:	1B8B3
	Nanodrop and 1000 µg/ml by Bradford method using BSA as the standard;			Dilutions recommandées:	WB 1:5000-1:50000 IHC 1:50-1:500
Hôte:	Mouse				
Isotype:	IgG1	MW calculé	1234 aa, 139 kDa		
Immunogen Catalog Number:	AG15845	MW observés:	150 kDa		

Applications

Applications testées:	IHC, WB, ELISA	Contrôles positifs:	
Demandes citées:	WB	WB :	cellules HepG2, cellules HEK-293, cellules HeLa, cellules HSC-T6, cellules Jurkat, cellules K-562, cellules LNCaP, cellules NIH/3T3, cellules Sp2/0, K-562, plaquettes de sang périphérique humain
Spécificité de l'espèce:	Humain, rat, souris	IHC :	tissu de cancer du sein humain, tissu de côlon humain
Espèces citées:	rat		
<i>Remarque-IHC: il est suggéré de démasquer l'antigène avec un tampon de TE buffer pH 9,0; (*) À défaut, 'le démasquage de l'antigène peut être effectué avec un tampon citrate pH 6,0.</i>			

Informations générales

PLCB3 is a member of the phosphoinositide phospholipase C beta enzyme family that catalyze the production of the secondary messengers diacylglycerol and inositol 1,4,5-triphosphate from phosphatidylinositol in G-protein-linked receptor-mediated signal transduction. Six subfamilies of PLCs (B, G, D, E, Z and H) constitute part of ubiquitous signaling cascades that translate hormonal signals into intracellular events, leading to alternations in cell function. PLCB isoforms 1-4 are stimulated by G-protein activation (Gαq/11 and/or Gβγ). Independent of its enzymatic activity, PLCB3 inhibits the proliferation of hematopoietic stem cells (HSCs) and myeloid cells.

Publications notables

Autrice	Pubmed ID	Journal	Application
Hongdong Song	35703476	J Food Sci	WB
Yijie Yang	32662128	J Food Biochem	WB
Wenna Zhang	31975557	J Cell Mol Med	WB

Stockage

Stockage:

Stocker à -20°C. Stable pendant un an après l'expédition.

Tampon de stockage:

PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3

L'aliquotage n'est pas nécessaire pour le stockage à -20C


*** Les 20ul contiennent 0,1% de BSA.

For technical support and original validation data for this product please contact:
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Données de validation sélectionnées




Various lysates were subjected to SDS PAGE followed by western blot with 66668-1-Ig (PLCB3 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Lamin B1 Monoclonal antibody (HRP-66095) as loading control.




Various lysates were subjected to SDS PAGE followed by western blot with 66668-1-Ig (PLCB3 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.




Various lysates were subjected to SDS PAGE followed by western blot with 66668-1-Ig (PLCB3 antibody) at dilution of 1:2500 incubated at room temperature for 1.5 hours.



Sp2/0 cells were subjected to SDS PAGE followed by western blot with 66668-1-Ig (PLCB3 antibody) at dilution of 1:2500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 66668-1-Ig (PLCB3 antibody) at dilution of 1:200 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 66668-1-Ig (PLCB3 antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).