

À des fins de recherche uniquement

Anticorps Monoclonal anti-TMEM173/STING



Numéro de catalogue: 66680-1-Ig

Phare

24 Publications

Informations de base

Numéro de catalogue: 66680-1-Ig	Numéro d'acquisition GenBank: BC047779	Méthode de purification: Purification par protéine A
Taille: 150ul, Concentration: 1900 µg/ml by Nanodrop and 1000 µg/ml by Bradford method using BSA as the standard;	Identification du gène (NCBI): 340061	CloneNo.: 1F1E1
Hôte: Mouse	Nom complet: transmembrane protein 173	Dilutions recommandées: WB 1:5000-1:50000 IHC 1:500-1:2000
Isotype: IgG2b	MW calculé: 379 aa, 42 kDa	
Immunogen Catalog Number: AG20363	MW observés: 37 kDa	

Applications

Applications testées:

IHC, WB, ELISA

Demandes citées:

IF, IHC, IP, WB

Spécificité de l'espèce:

Humain, rat

Espèces citées:

Humain, souris

Remarque-IHC: il est suggéré de démasquer l'antigène avec un tampon de TE buffer pH 9,0; (*) A défaut, 'le démasquage de l'antigène peut être effectué avec un tampon citrate pH 6,0.

Contrôles positifs:

WB : cellules HeLa, cellules HEK-293, cellules HepG2, cellules HSC-T6, cellules Jurkat, cellules MOLT-4, cellules THP-1

IHC : tissu d'amygdalite humain, tissu splénique humain

Informations générales

Stimulator of interferon genes (STING, also known as ERIS, MITA and MPYS, and encoded by TMEM173) is a transmembrane adaptor protein that facilitates innate immune signaling (PMID: 18724357). STING is widely expressed in various cell types such as endothelial cells, epithelial cells, T cells, macrophages, and dendritic cells (PMID: 26603901). It is predominantly located in the endoplasmic reticulum (ER). STING functions as a sensor of cytosolic DNA and promotes the production of type I interferons and pro-inflammatory cytokines. STING is a 379 amino acid protein with a calculated molecular weight of 42 kDa. It has been observed at 35-40 kDa (PMID: 27324217; 29632140; 30918080), and 70-80 kDa corresponding to the expected size of a STING dimer (PMID: 25790474; 29491158).

Publications notables

Autrice	Pubmed ID	Journal	Application
Mu-zi Li	36238763	J Inflamm Res	WB,IHC,IF
Jiaqi Shi	34182334	Cell Immunol	WB
Ming Zou	35347235	Cell Death Differ	WB

Stockage

Stockage:

Stocker à -20°C. Stable pendant un an après l'expédition.

Tampon de stockage:

PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3

L'aliquotage n'est pas nécessaire pour le stockage à -20C

*** Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Données de validation sélectionnées




Various lysates were subjected to SDS PAGE followed by western blot with 66680-1-Ig (TMEM173/STING antibody) at dilution of 1:15000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Lamin B1 Monoclonal antibody (HRP-66095) as loading control.



WB result of TMEM173/STING antibody (66680-1-Ig; 1:1200; incubated at room temperature for 1.5 hours) with sh-Control and sh-TMEM173/STING transfected THP-1 cells.



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 66680-1-Ig (TMEM173/STING antibody) at dilution of 1:1000 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 66680-1-Ig (TMEM173/STING antibody) at dilution of 1:1000 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).