

À des fins de recherche uniquement

# Anticorps Monoclonal anti-Cytokeratin 13

Numéro de catalogue: 66684-1-Ig



## Informations de base

Numéro de catalogue:	BC002661	Méthode de purification:
66684-1-Ig		Purification par protéine G
Taille:	Identification du gène (NCBI):	CloneNo.:
150ul , Concentration: 1500 µg/ml by Nanodrop and 1000 µg/ml by Bradford method using BSA as the standard;	3860 Nom complet: keratin 13	1E10G2
Hôte:	MW calculé	Dilutions recommandées:
Mouse	50 kDa	WB 1:5000-1:50000 IHC 1:1000-1:4000 IF 1:200-1:800
Isotype:	MW observés:	
IgG1	50 kDa	
Immunogen Catalog Number:		
AG0217		

## Applications

Applications testées:	Contrôles positifs:
FC, IF, IHC, WB, ELISA	WB: cellules A431, tissu cutané de rat
Spécificité de l'espèce:	IHC : tissu de cancer du col de l'utérus humain, tissu d'amygdalite humain, tissu de cancer de l'œsophage humain
Humain, rat	IF : tissu de cancer du col de l'utérus humain,
<b>Remarque-IHC: il est suggéré de démasquer l'antigène avec un tampon de TE buffer pH 9,0; (*) À défaut, le démasquage de l'antigène peut être effectué avec un tampon citrate pH 6,0.</b>	

## Informations générales

Keratins are a large family of proteins that form the intermediate filament cytoskeleton of epithelial cells, which are classified into two major sequence types. Type I keratins are a group of acidic intermediate filament proteins, including K9-K23, and the hair keratins Ha1-Ha8. Type II keratins are the basic or neutral counterparts to the acidic type I keratins, including K1-K8, and the hair keratins, Hb1-Hb6. This type I cytokeratin is paired with keratin 4 and expressed in the suprabasal layers of non-cornified stratified epithelia. The Keratin 13 (KRT13) gene encodes a type I acidic keratin which is expressed in the differentiated cells of non-cornified stratified squamous epithelia. This antibody can react with Cytokeratin 15.

## Stockage

**Stockage:**  
Stocker à -20°C. Stable pendant un an après l'expédition.  
**Tampon de stockage:**  
PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3  
L'aliquotage n'est pas nécessaire pour le stockage à -20C




\*\*\* Les 20ul contiennent 0,1% de BSA.

For technical support and original validation data for this product please contact:  
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free  
in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com  
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Données de validation sélectionnées




A431 cells were subjected to SDS PAGE followed by western blot with 66684-1-Ig (KRT13 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.

Immunohistochemical analysis of paraffin-embedded human cervical cancer tissue slide using 66684-1-Ig (KRT13 antibody) at dilution of 1:2000 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).

Immunohistochemical analysis of paraffin-embedded human cervical cancer tissue slide using 66684-1-Ig (KRT13 antibody) at dilution of 1:2000 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed human cervical cancer tissue using Cytokeratin 13 antibody (66684-1-Ig, Clone: 1E10G2) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



1X10<sup>6</sup> A431 cells were intracellularly stained with 0.2 ug Anti-Human Cytokeratin 13 (66684-1-Ig, Clone:1E10G2) and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (green), and 0.2 ug Control Antibody. Cells were fixed with 90% MeOH.