

À des fins de recherche uniquement

# Anticorps Monoclonal anti-CNKSR1

Numéro de catalogue: 66796-1-Ig



## Informations de base

Numéro de catalogue: 66796-1-Ig	Numéro d'acquisition GenBank: BC011604	Méthode de purification: Purification par protéine G
Taille: 150ul, Concentration: 1700 µg/ml by Nanodrop and 1000 µg/ml by Bradford method using BSA as the standard;	Identification du gène (NCBI): 10256	CloneNo.: 1E11H8
Hôte: Mouse	Nom complet: connector enhancer of kinase suppressor of Ras 1	Dilutions recommandées: WB 1:1000-1:4000
Isotype: IgG1	MW calculé: 80 kDa	
Immunogen Catalog Number: AG27855	MW observés: 90 kDa	

## Applications

Applications testées: WB, ELISA	Contrôles positifs: WB : cellules MCF-7, cellules Caco-2, cellules HEK-293, cellules HT-29
Spécificité de l'espèce: Humain	

## Informations générales

CNKSR1 is a protein containing several motifs involved in protein-protein interaction, including PDZ, PH (Pleckstrin homology), and SAM (sterile alpha motif) domains. The encoded protein acts as a scaffold component for receptor tyrosine kinase signaling and may mediate crosstalk between different signaling pathways. CNKSR1 is required in multiple receptor tyrosine kinase pathways where it appears to be a tyrosine phosphorylation target.

## Stockage

Stockage:  
Stocker à -20°C. Stable pendant un an après l'expédition.  
Tampon de stockage:  
PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3  
L'aliquotage n'est pas nécessaire pour le stockage à -20C

\*\*\* Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free  
in USA), or 1(312) 455-8498 (outside USA)

E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Données de validation sélectionnées



MCF-7 cells were subjected to SDS PAGE followed by western blot with 66796-1-Ig (CNKSR1 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.