

À des fins de recherche uniquement

Anticorps Monoclonal anti-FRZB

Numéro de catalogue: 67081-1-Ig **1 Publications**



Informations de base

Numéro de catalogue: 67081-1-Ig	Numéro d'acquisition GenBank: BC027855	Méthode de purification: Purification par protéine G
Taille: 150ul , Concentration: 2200 µg/ml by Nanodrop and 1000 µg/ml by Bradford method using BSA as the standard;	Identification du gène (NCBI): 2487	CloneNo.: 2B4A1
Hôte: Mouse	Nom complet: frizzled-related protein	Dilutions recommandées: WB 1:5000-1:50000
Isotype: IgG1	MW calculé: 325 aa, 36 kDa	
Immunogen Catalog Number: AG28147	MW observés: 36 kDa	

Applications

Applications testées: WB, ELISA	Contrôles positifs: WB : cellules LNCaP, cellules 4T1, cellules MKN-45, cellules SGC-7901
Demandes citées: WB	
Spécificité de l'espèce: Humain, souris	
Espèces citées: Humain	

Informations générales

Frzb (also known as sFRP3) is a member of the secreted frizzled related protein (SFRP) family, originally identified as a chondrogenic factor during bone morphogenesis (PMID: 29209231). It was subsequently shown to be a regulator in Wnt signaling, and also have a role in regulating cell growth and differentiation in specific cell types. FRZB was seen to be acting as an oncogene in some cancer, such as metastatic renal cancer.

Publications notables

Autrice	Pubmed ID	Journal	Application
Xiaoping Hu	33789447	Stem Cells Dev	WB

Stockage


Stockage:
Stocker à -20 °C
Tampon de stockage:
PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3
L'aliquotage n'est pas nécessaire pour le stockage à -20C

*** Les 20ul contiennent 0,1% de BSA.

For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



Various lysates were subjected to SDS PAGE followed by western blot with 67081-1-Ig (FRZB antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.