

À des fins de recherche uniquement

Anticorps Monoclonal anti-DNAJA1

Numéro de catalogue: 67184-1-Ig



Informations de base

Numéro de catalogue: 67184-1-Ig	Numéro d'acquisition GenBank: BC008182	Méthode de purification: Précipitation des euglobulines
Taille: 150ul , Concentration: 500 µg/ml by Bradford method using BSA as the standard;	Identification du gène (NCBI): 3301	CloneNo.: 1B10A10
Hôte: Mouse	Nom complet: DnaJ (Hsp40) homolog, subfamily A, member 1	Dilutions recommandées: WB 1:1000-1:6000 IF 1:200-1:800
Isotype: IgM	MW calculé 397 aa, 45 kDa	
Immunogen Catalog Number: AG2284	MW observés: 45 kDa	

Applications

Applications testées:

IF, WB, ELISA

Spécificité de l'espèce:

Humain, rat, souris

Contrôles positifs:

WB : cellules A549, cellules HEK-293, cellules HeLa,
cellules HepG2, cellules HSC-T6, cellules Jurkat,
cellules K-562, cellules MCF-7, cellules NIH/3T3

IF : cellules HepG2,

Informations générales

DNAJA1 is a member of the DnaJ proteins (also known as Hsp40 or Hsc40) which are cochaperones to DnaK (Hsp70) and function to modulate protein assembly, disassembly, and translocation. DNAJA1 has been reported to get involved in various processes, including virus infection, tau clearance, and mutant p53 degradation.

Stockage

Stockage:

Stocker à -20°C. Stable pendant un an après l'expédition.

Tampon de stockage:

PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3

L'aliquotage n'est pas nécessaire pour le stockage à -20C

***** Les 20ul contiennent 0,1% de BSA.**


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



Various lysates were subjected to SDS PAGE followed by western blot with 67184-1-Ig (DNAJA1 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using DNAJA1 antibody (67184-1-Ig, Clone: 1B10A10) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).