

À des fins de recherche uniquement

Anticorps Monoclonal anti-Septin 9

Numéro de catalogue: 67548-1-Ig



Informations de base

Numéro de catalogue: 67548-1-Ig	Numéro d'acquisition GenBank: BC021192	Méthode de purification: Purification par protéine G
Taille: 150ul , Concentration: 1000 µg/ml by Nanodrop and 500 µg/ml by Bradford method using BSA as the standard;	Identification du gène (NCBI): 10801	CloneNo.: 1C6F10
Hôte: Mouse	Nom complet: septin 9	Dilutions recommandées: WB 1:20000-1:100000
Isotype: IgG1	MW calculé 64 kDa	
Immunogen Catalog Number: AG18047	MW observés: 42 kDa	

Applications

Applications testées: WB, ELISA	Contrôles positifs:
Spécificité de l'espèce: Humain, porc, rat	WB : tissu de côlon de porc, cellules A549, cellules HeLa, cellules HepG2, cellules MCF-7, tissu de côlon de rat

Informations générales

Septin 9 belongs to the septin family which is a group of highly conserved GTP-binding proteins in eukaryotes. SEPT9 gene methylation has been implicated as a biomarker for colorectal cancer(CRC). this antibody can identify all of the isoforms about SEPT9.

Stockage

Stockage:
Stocker à -20°C. Stable pendant un an après l'expédition.
Tampon de stockage:
PBS avec azoture de sodium à 0,1 % et glycérol à 50 % pH 7,3
L'aliquotage n'est pas nécessaire pour le stockage à -20C

***** Les 20ul contiennent 0,1% de BSA.**


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Données de validation sélectionnées



Various lysates were subjected to SDS PAGE followed by western blot with 67548-1-Ig (Septin 9 antibody) at dilution of 1:50000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 67548-1-Ig (Septin 9 antibody) at dilution of 1:50000 incubated at room temperature for 1.5 hours.



rat colon tissue were subjected to SDS PAGE followed by western blot with 67548-1-Ig (Septin 9 antibody) at dilution of 1:50000 incubated at room temperature for 1.5 hours.