

À des fins de recherche uniquement

Anticorps Monoclonal anti-GSTT1

Numéro de catalogue: 67645-1-Ig



Informations de base

Numéro de catalogue:	BC007065	Méthode de purification:
67645-1-Ig	Purification par protéine A	
Taille:	Identification du gène (NCBI):	CloneNo.:
150ul , Concentration: 2200 µg/ml by 2952	Nom complet:	1A4C11
Nanodrop and 1000 µg/ml by Bradford method using BSA as the standard;	glutathione S-transferase theta 1	Dilutions recommandées:
Hôte:	MW calculé	WB 1:2000-1:10000
Mouse	240 aa, 27 kDa	IHC 1:250-1:1000
Isotype:	MW observés:	IF 1:50-1:500
IgG2b	28 kDa, 30 kDa	
Immunogen Catalog Number:		
AG8573		

Applications

Applications testées:	Contrôles positifs:
IF, IHC, WB, ELISA	WB : tissu pulmonaire de porc, cellules K-562, tissu cérébral de souris, tissu hépatique de porc, tissu hépatique de rat
Spécificité de l'espèce:	IHC : tissu hépatique de souris,
Humain	IF : tissu hépatique de souris,
Remarque-IHC: il est suggéré de démasquer l'antigène avec un tampon de TE buffer pH 9,0; (*) À défaut, le démasquage de l'antigène peut être effectué avec un tampon citrate pH 6,0.	

Informations générales

GSTT1 belongs to the GST superfamily and Theta family. It is a conjugation of reduced glutathione to a wide number of exogenous and endogenous hydrophobic electrophiles. GSTT1 acts on 1,2-epoxy-3-(4-nitrophenoxyl)propane, phenethylisothiocyanate 4-nitrobenzyl chloride and 4-nitrophenethyl bromide. It displays glutathione peroxidase activity with cumene hydroperoxide. GSTM1 and GSTT1 has been suggested as a risk factor in various cancers. GSTT1 is important in the detoxification of unidentified xenobiotics in the large intestine.

Stockage


Stockage:
Stocker à -20 °C.
Tampon de stockage:
PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3
L'aliquotage n'est pas nécessaire pour le stockage à -20°C

*** Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Données de validation sélectionnées




Immunohistochemical analysis of paraffin-embedded mouse liver tissue slide using 67645-1 Ig (GSTT1 antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Various lysates were subjected to SDS PAGE followed by western blot with 67645-1-Ig (GSTT1 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded mouse liver tissue slide using 67645-1-Ig (GSTT1 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed mouse liver tissue using GSTT1 antibody (67645-1 Ig, Clone: 1A4C11) at dilution of 1:200 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).