

À des fins de recherche uniquement

Anticorps Monoclonal anti-AATF

Numéro de catalogue: 67868-1-Ig **2 Publications**



Informations de base

Numéro de catalogue: 67868-1-Ig	Numéro d'acquisition GenBank: BC000591	Méthode de purification: Purification par protéine G
Taille: 150ul, Concentration: 1000 µg/ml by Nanodrop;	Identification du gène (NCBI): 26574	CloneNo.: 2A6G6
Hôte: Mouse	Nom complet: apoptosis antagonizing transcription factor	Dilutions recommandées: WB 1:5000-1:50000
Isotype: IgG1	MW calculé 63 kDa	
Immunogen Catalog Number: AG0195	MW observés: 80-90 kDa	

Applications

Applications testées:
FC, WB, ELISA

Demandes citées:
IHC, WB

Spécificité de l'espèce:
Humain, rat, souris

Espèces citées:
Humain

Contrôles positifs:

WB : cellules HCT 116, cellules HeLa, cellules HepG2,
cellules Jurkat, cellules K-562, cellules LNCaP, cellules
Neuro-2a

Informations générales

Publications notables

Autrice	Pubmed ID	Journal	Application
Lin Fu	34785906	Onco Targets Ther	WB,IHC
Shutao Tan	35002255	Onco Targets Ther	WB,IHC

Stockage

Stockage:

Stocker à -20°C. Stable pendant un an après l'expédition.

Tampon de stockage:

PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3

L'aliquotage n'est pas nécessaire pour le stockage à -20C

***** Les 20ul contiennent 0,1% de BSA.**


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Données de validation sélectionnées



Various lysates were subjected to SDS PAGE followed by western blot with 67868-1-Ig (AATF antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Neuro-2a cells were subjected to SDS PAGE followed by western blot with 67868-1-Ig (AATF antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



1X10⁶ HeLa cells were intracellularly stained with 0.5 ug Anti-Human AATF (67868-1-Ig, Clone:2A6G6) (red) labeled with FlexAble CoraLite® Plus 555 Antibody Labeling Kit for Mouse IgG1 (KFA022), or 0.5 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).