

À des fins de recherche uniquement

Anticorps Monoclonal anti-PLEKHB2

Numéro de catalogue: 67877-1-Ig



Informations de base

Numéro de catalogue: 67877-1-Ig	Numéro d'acquisition GenBank: BC013991	Méthode de purification: Purification par protéine G
Taille: 150ul, Concentration: 1000 µg/ml by Nanodrop;	Identification du gène (NCBI): 55041	CloneNo.: 1H9D7
Hôte: Mouse	Nom complet: pleckstrin homology domain containing, family B (evectins) member 2	Dilutions recommandées: WB 1:2000-1:10000
Isotype: IgG1	MW calculé 222 aa, 25 kDa	
Immunogen Catalog Number: AG9231	MW observés: 19-25;30-34 kDa	

Applications

Applications testées: WB, ELISA	Contrôles positifs:
Spécificité de l'espèce: Humain	WB : cellules Caco-2, cellules BxPC-3, cellules Colo320, cellules HL-60, cellules HT-29, cellules MG U- 87, cellules NCCIT, cellules SW480, tissu de côlon de porc

Informations générales

PLEKHB2, also named as EVT2 (Evectin-2), is involved in retrograde transport of recycling endosomes. PLEKHB2 has been identified as post-Golgi proteins of unknown function that have a PH domain that typically binds polyphosphoinositides. PLEKHB2 is expressed in a broad range of tissues. PLEKHB2 has some isoforms with MW 19-26 kDa and 34 kDa. This antibody recognizes all the isoforms of PLEKHB2.

Stockage

Stockage:
Stocker à -20°C. Stable pendant un an après l'expédition.
Tampon de stockage:
PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3
L'aliquotage n'est pas nécessaire pour le stockage à -20C

*** Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



Various lysates were subjected to SDS PAGE followed by western blot with 67877-1-Ig (PLEKHB2 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control.