

À des fins de recherche uniquement

Anticorps Monoclonal anti-GMEB2

Numéro de catalogue: 68010-1-Ig

Phare



Informations de base

Numéro de catalogue:	68010-1-Ig	Numéro d'acquisition GenBank:	BC036305	Méthode de purification:	Purification par protéine G
Taille:	150ul, Concentration: 1000 µg/ml by Nanodrop;	Identification du gène (NCBI):	26205	CloneNo.:	2H10H8
Hôte:	Mouse	Nom complet:	glucocorticoid modulatory element binding protein 2	Dilutions recommandées:	WB 1:2000-1:10000
Isotype:	IgG1	MW calculé	530 aa, 56 kDa		
Immunogen Catalog Number:	AG31393	MW observés:	68 kDa		

Applications

Applications testées:	WB, ELISA	Contrôles positifs:	WB : cellules HeLa, cellules HEK-293, cellules HSC-T6, cellules Jurkat, cellules K-562, cellules MCF-7, cellules NIH/3T3, cellules PC-12, cellules RAW 264.7
Spécificité de l'espèce:	Humain, rat, souris		

Informations générales

GMEB-2 (glucocorticoid modulatory element-binding protein 2), also known as PIF79 (Parvovirus initiation factor p79) or P79PIF, is a DNA-binding protein that plays a crucial role in modulating transcription upon activation by steroid hormones. GMEB-2 is ubiquitously expressed with preferential expression in developmentally important tissues.

Stockage


Stockage:
Stocker à -20°C. Stable pendant un an après l'expédition.
Tampon de stockage:
PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3
L'aliquotage n'est pas nécessaire pour le stockage à -20C

*** Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:
T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)
E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Données de validation sélectionnées




WB result of GMEB2 antibody (68010-1-Ig; 1:4000; incubated at room temperature for 1.5 hours) with sh-Control and sh-GMEB2 transfected MCF-7 cells.



HeLa cells were subjected to SDS PAGE followed by western blot with 68010-1-Ig (GMEB2 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Beta Actin Monoclonal antibody (HRP-66009) as loading control.



Various lysates were subjected to SDS PAGE followed by western blot with 68010-1-Ig (GMEB2 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Beta Actin Monoclonal antibody (HRP-66009) as loading control.



Various lysates were subjected to SDS PAGE followed by western blot with 68010-1-Ig (GMEB2 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.