

À des fins de recherche uniquement

Anticorps Monoclonal anti-ACOT7

Numéro de catalogue: 68140-1-Ig



Informations de base

Numéro de catalogue:	BC017365	Méthode de purification:
68140-1-Ig		Purification par protéine A
Taille:	Identification du gène (NCBI):	CloneNo.:
150ul , Concentration: 1000 µg/ml by Nanodrop;	11332	1E9C7
Hôte:	Nom complet:	Dilutions recommandées:
Mouse	acyl-CoA thioesterase 7	WB 1:5000-1:50000
Isotype:	MW calculé	
IgG2a	370 aa, 40 kDa	
Immunogen Catalog Number:	MW observés:	
AG8975	37 kDa	

Applications

Applications testées:	WB, ELISA	Contrôles positifs:
Spécificité de l'espèce:	Humain, Lapin, porc, rat, souris	WB : cellules LNCaP, cellules HEK-293, cellules HeLa, cellules HSC-T6, cellules Jurkat, cellules K-562, cellules NIH/3T3, tissu cérébral de lapin, tissu cérébral de porc

Informations générales

Acyl-CoA thioesterases (ACOTs) are enzymes that catalyze hydrolysis of fatty acyl-CoAs to free fatty acids and CoA-SH. ACOTs are expressed ubiquitously in prokaryotes and eukaryotes (PMID: 11755680). ACOT7 also known as BACH, CTE-II, ACT, ACH1, BACHa, is one of the most extensively studied ACOTs as a major isoform of the ACOT family (PMID: 17563367). The enzyme is localized to the cytosol and is highly expressed in brain tissue and testis tissue, showing a preference for long-chain acyl-CoA substrates with fatty acid chains of 8-16 carbon atoms (C8-C16) (PMID: 28518146). It has been identified that ACOT7 plays a role in neurotoxic prevention and is a candidate drug target in inflammatory disease (PMID: 17563367, 23459938, 10509865). ACOT7 has 7 isoforms as molecular between 27-40 kDa, it may also form dimer (~70 kDa) (PMID: 20470824).

Stockage

Stockage:
Stocker à -20°C. Stable pendant un an après l'expédition.

Tampon de stockage:
PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3
L'aliquotage n'est pas nécessaire pour le stockage à -20°C


*** Les 20ul contiennent 0,1% de BSA.

For technical support and original validation data for this product please contact:
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



Various lysates were subjected to SDS PAGE followed by western blot with 68140-1-Ig (ACOT7 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Alpha Tubulin Monoclonal antibody (HRP-66031) as loading control.