

À des fins de recherche uniquement

Anticorps Monoclonal anti-AIDA

Numéro de catalogue: 68153-1-Ig



Informations de base

Numéro de catalogue: 68153-1-Ig	Numéro d'acquisition GenBank: BC015535	Méthode de purification: Purification par protéine G
Taille: 150ul, Concentration: 1000 µg/ml by Nanodrop;	Identification du gène (NCBI): 64853	CloneNo.: 1D11A1
Hôte: Mouse	Nom complet: axin interactor, dorsalization associated	Dilutions recommandées: WB 1:2000-1:10000 IF 1:400-1:1600
Isotype: IgG1	MW calculé 306 aa, 35 kDa	
Immunogen Catalog Number: AG20597	MW observés: 35 kDa	

Applications

Applications testées:
IF, WB, ELISA
Spécificité de l'espèce:
Humain

Contrôles positifs:

WB : cellules HeLa, cellules COLO 320, cellules HepG2,
cellules HT-29, cellules K-562, cellules U2OS
IF : cellules A431,

Informations générales

AIDA, also named as C1orf80, acts as a ventralizing factor during embryogenesis. It inhibits axin-mediated JNK activation by binding axin and disrupting axin homodimerization. It's highest expression in the heart and skeletal muscle. AIDA has three isoforms with MW 35 kDa, 16 kDa and 26 kDa.

Stockage

Stockage:
Stocker à -20°C. Stable pendant un an après l'expédition.
Tampon de stockage:
PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3
L'aliquotage n'est pas nécessaire pour le stockage à -20C

*** Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Données de validation sélectionnées




Various lysates were subjected to SDS PAGE followed by western blot with 68153-1-Ig (AIDA antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 68153-1-Ig (AIDA antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



HepG2 cells were subjected to SDS PAGE followed by western blot with 68153-1-Ig (AIDA antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Methanol) fixed A431 cells using AIDA antibody (68153-1-Ig, Clone: 1D11A1) at dilution of 1:800 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).