

À des fins de recherche uniquement

Anticorps Monoclonal anti-TOLLIP

Numéro de catalogue: 68170-1-Ig Phare



Informations de base

Numéro de catalogue:	68170-1-Ig	Numéro d'acquisition GenBank:	BC012057	Méthode de purification:	Purification par protéine A
Taille:	150ul , Concentration: 1000 µg/ml by Nanodrop;	Identification du gène (NCBI):	54472	CloneNo.:	1B3C4
Hôte:	Mouse	Nom complet:	toll interacting protein	Dilutions recommandées:	WB 1:5000-1:50000 IHC 1:400-1:1600 IF 1:400-1:1600
Isotype:	IgG1	MW calculé	30 kDa		
Immunogen Catalog Number:	AG32731	MW observés:	30 kDa		

Applications

Applications testées:	FC, IF, IHC, WB, ELISA	Contrôles positifs:	
Spécificité de l'espèce:	Humain, rat, souris	WB :	tissu cérébral de porc, cellules HepG2, cellules Jurkat, cellules K-562, cellules LNCaP, plaquettes de sang périphérique humain, tissu cérébral de lapin, tissu cérébral de rat, tissu cérébral de souris
Remarque-IHC: il est suggéré de démasquer l'antigène avec un tampon de TE buffer pH 9.0; (*) À défaut, le démasquage de l'antigène peut être effectué avec un tampon citrate pH 6.0.		IHC :	tissu de côlon de souris, tissu de côlon de rat
		IF :	cellules U2OS,

Informations générales

TOLLIP (Toll interacting protein), is an inhibitory adaptor protein. Tollip is a negative regulator of IL-1R and Toll-like receptors signalling pathway, involved in inflammatory immune response and inhibiting IRAK1 phosphorylation and kinase activity (PMID:11441107, PMID:10854325, PMID:11751856). The TOLLIP-dependent selective autophagy pathway plays a role in clearance of cytotoxic polyQ protein aggregates (PubMed:25042851).

Stockage

Stockage:

Stocker à -20°C. Stable pendant un an après l'expédition.

Tampon de stockage:

PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3


L'aliquotage n'est pas nécessaire pour le stockage à -20°C

*** Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Données de validation sélectionnées







Various lysates were subjected to SDS PAGE followed by western blot with 68170-1-Ig (TOLLIP antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




Various lysates were subjected to SDS PAGE followed by western blot with 68170-1-Ig (TOLLIP antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Beta Actin Monoclonal antibody (HRP-66009) as loading control.



WB result of TOLLIP antibody (68170-1-Ig; 1:1000; incubated at room temperature for 1.5 hours) with sh-Control and sh-TOLLIP transfected HepG2 cells.



Immunofluorescent analysis of (-20°C Methanol) fixed U2OS cells using TOLLIP antibody (68170-1-Ig, Clone: 1B3C4) at dilution of 1:800 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



1×10^6 Jurkat cells were intracellularly stained with 0.4 ug Anti-Human TOLLIP (68170-1-Ig, Clone:1B3C4) and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Mouse IgG1 Isotype Control (MOPC-21) (65124-1-Ig, Clone: MOPC-21) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).