

À des fins de recherche uniquement

Anticorps Monoclonal anti-NOTCH2/NOTCH2NL



Numéro de catalogue: 68212-1-Ig

Informations de base

Numéro de catalogue: 68212-1-Ig	Numéro d'acquisition GenBank: BC019835	Méthode de purification: Purification par protéine G
Taille: 150ul , Concentration: 1000 µg/ml by Nanodrop;	Identification du gène (NCBI): 388677	CloneNo.: 2D5E1
Hôte: Mouse	Nom complet: Notch homolog 2 (Drosophila) N- terminal like	Dilutions recommandées: WB 1:5000-1:50000
Isotype: IgG1	MW calculé 265 kDa	
Immunogen Catalog Number: AG1679	MW observés: 250-265 kDa	

Applications

Applications testées: WB, ELISA	Contrôles positifs: WB : cellules hTERT-RPE1, cellules MG U-87
Spécificité de l'espèce: Humain	

Informations générales

Notch 2 is a member of the notch family. Members of this Type 1 transmembrane protein family share structural characteristics including an extracellular domain consisting of multiple epidermal growth factor-like (EGF) repeats, and an intracellular domain consisting of multiple, different domain types. Notch family members play a role in a variety of developmental processes by controlling cell fate decisions. Notch2, like other Notch genes, functions in development of various tissues. This antibody can recognize NOTCH2 and NOTCH2NL. The antibody detects the full-length NOTCH2 protein about 250-265 kDa.

Stockage

Stockage:
Stocker à -20°C. Stable pendant un an après l'expédition.
Tampon de stockage:
PBS avec azote de sodium à 0,02 % et glycérol à 50 % pH 7,3
L'aliquotage n'est pas nécessaire pour le stockage à -20C

***** Les 20ul contiennent 0,1% de BSA.**

For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



hTERT-RPE1 cells were subjected to SDS PAGE followed by western blot with 68212-1-Ig (NOTCH2 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.