

À des fins de recherche uniquement

Anticorps Monoclonal anti-HIGD1A

Numéro de catalogue: 68231-1-Ig



Informations de base

Numéro de catalogue: 68231-1-Ig	Numéro d'acquisition GenBank: BC000601	Méthode de purification: Purification par protéine A
Taille: 150ul, Concentration: 1000 µg/ml by Nanodrop;	Identification du gène (NCBI): 25994	CloneNo.: 3A8E2
Hôte: Mouse	Nom complet: HIG1 hypoxia inducible domain family, member 1A	Dilutions recommandées: WB 1:1000-1:6000 IF 1:200-1:800
Isotype: IgG2b	MW calculé 93 aa, 10 kDa	
Immunogen Catalog Number: AG14027	MW observés: 10 kDa	

Applications

Applications testées:
IF, WB, ELISA

Spécificité de l'espèce:
Humain

Contrôles positifs:

WB : cellules LNCaP, cellules HeLa, cellules HepG2, cellules K-562, tissu placentaire humain

IF : cellules U2OS,

Informations générales

HIG1 domain family member 1A (HIGD1A) is a proposed subunit of cytochrome c oxidase (COX, complex IV), which is the terminal component of the mitochondrial respiratory chain that catalyzes the reduction of oxygen to water. May play a role in the assembly of respiratory supercomplexes (PMID:22342701). It is induced by hypoxia induction.

Stockage

Stockage:
Stocker à -20°C. Stable pendant un an après l'expédition.

Tampon de stockage:
PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3
L'aliquotage n'est pas nécessaire pour le stockage à -20C

***** Les 20ul contiennent 0,1% de BSA.**


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



Immunofluorescent analysis of (4% PFA) fixed U2OS cells using HIGD1A antibody (68231-1-Ig, Clone: 3A8E2) at dilution of 1:400 and CoraLite@488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L), CL594-Phalloidin (red).



Various lysates were subjected to SDS PAGE followed by western blot with 68231-1-Ig (HIGD1A antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control.