

À des fins de recherche uniquement

# Anticorps Monoclonal anti-SNX17

Numéro de catalogue: 68256-1-Ig



## Informations de base

Numéro de catalogue:	68256-1-Ig	Numéro d'acquisition GenBank:	BC002524	Méthode de purification:	Purification par protéine G
Taille:	150ul, Concentration: 500 µg/ml by Nanodrop;	Identification du gène (NCBI):	9784	CloneNo.:	3B8F6
Hôte:	Mouse	Nom complet:	sorting nexin 17	Dilutions recommandées:	WB 1:5000-1:50000 IF 1:200-1:800
Isotype:	IgG1	MW calculé:	53 kDa		
Immunogen Catalog Number:	AG32624	MW observés:	55 kDa		

## Applications

Applications testées:  
IF, WB, ELISA

Spécificité de l'espèce:  
Humain, porc, rat

Contrôles positifs:

WB : cellules LNCaP, cellules HeLa, cellules Jurkat, tissu cérébral de porc, tissu cérébral de rat

IF : cellules HepG2, cellules HeLa

## Informations générales

Sorting nexin 17 (SNX17, synonym: KIAA0064) is a member of the sorting nexin family. Members of this family contain a phox (PX) domain, which is a phosphoinositide binding domain, and are involved in intracellular trafficking. SNX17 does not contain a coiled coil region, like some family members, but contains a B41 domain. This protein interacts with the cytoplasmic domain of P-selectin, and may function in the intracellular trafficking of P-selectin.

## Stockage

Stockage:

Stocker à -20°C. Stable pendant un an après l'expédition.

Tampon de stockage:

PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3

L'aliquotage n'est pas nécessaire pour le stockage à -20C

\*\*\* Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Données de validation sélectionnées



Various lysates were subjected to SDS PAGE followed by western blot with 68256-1-Ig (SNX17 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using SNX17 antibody (68256-1-Ig, Clone: 3B8F6) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L), CL594-Phalloidin (red).