

À des fins de recherche uniquement

Anticorps Monoclonal anti-DHX32

Numéro de catalogue: 68306-1-Ig



Informations de base

Numéro de catalogue: 68306-1-Ig	Numéro d'acquisition GenBank: BC002473	Méthode de purification: Purification par protéine A
Taille: 150ul , Concentration: 1000 µg/ml by Nanodrop;	Identification du gène (NCBI): 55760	CloneNo.: 2A4G9
Hôte: Mouse	Nom complet: DEAH (Asp-Glu-Ala-His) box polypeptide 32	Dilutions recommandées: WB 1:2000-1:10000
Isotype: IgG2b	MW calculé: 743 aa, 84 kDa	
Immunogen Catalog Number: AG13956	MW observés: 84 kDa	

Applications

Applications testées:
WB, ELISA

Spécificité de l'espèce:
Humain, rat, souris

Contrôles positifs:

WB : cellules A431, cellules A549, cellules HeLa, cellules K-562, cellules MCF-7, cellules U2OS, tissu testiculaire de rat, tissu testiculaire de souris

Informations générales

DHX32, an RNA helicase, belongs to the DEAH box RNA helicase family by virtue of its homology to the conserved helicase domain. DHX32 is abundantly expressed in human and murine models, participating in cell transcription and differentiation, and being capable of regulating T-cell apoptosis. DHX32 is also associated with some malignant tumors, such as lymphoma, colorectal cancer, and breast cancer (PMID: 35117530). DHX32 has 2 isoforms with molecular weights of 75 and 84 kDa, respectively.

Stockage

Stockage:
Stocker à -20°C. Stable pendant un an après l'expédition.
Tampon de stockage:
PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3
L'aliquotage n'est pas nécessaire pour le stockage à -20C

***** Les 20ul contiennent 0,1% de BSA.**


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



Various lysates were subjected to SDS PAGE followed by western blot with 68306-1-Ig (DHX32 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.