

À des fins de recherche uniquement

Anticorps Monoclonal anti-AZI2

Numéro de catalogue: 68326-1-Ig



Informations de base

Numéro de catalogue: 68326-1-Ig	Numéro d'acquisition GenBank: BC001139	Méthode de purification: Purification par protéine G
Taille: 150ul , Concentration: 1000 µg/ml by Nanodrop;	Identification du gène (NCBI): 64343	CloneNo.: 2F1G5
Hôte: Mouse	Nom complet: 5-azacytidine induced 2	Dilutions recommandées: WB 1:5000-1:50000 IF 1:500-1:2000
Isotype: IgG1	MW calculé 45 kDa	
Immunogen Catalog Number: AG6755	MW observés: 45 kDa	

Applications

Applications testées:
IF, WB, ELISA

Spécificité de l'espèce:
Humain

Contrôles positifs:

WB : cellules U2OS, cellules 4T1, cellules HEK-293, cellules HeLa, cellules HSC-T6, cellules K-562, cellules LNCaP, cellules MDA-MB-231

IF : cellules HeLa,

Informations générales

AZI2, also known as NAP1, is a TNF receptor (TNFR)-associated factor family member-associated NF-κB activator-binding kinase 1-binding protein that regulates the production of IFNs. It is an adapter protein which binds TBK1 and IKKε playing a role in antiviral innate immunity. NAP1 is a kind of adapter protein which binds TBK1 and IKKε playing a role in antiviral innate immunity. (PMID: 21931631)

Stockage

Stockage:
Stocker à -20°C. Stable pendant un an après l'expédition.

Tampon de stockage:
PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3

L'aliquotage n'est pas nécessaire pour le stockage à -20C

*** Les 20ul contiennent 0,1% de BSA.



For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



Various lysates were subjected to SDS PAGE followed by western blot with 68326-1-Ig (AZI2 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control.

Immunofluorescent analysis of (-20°C Methanol) fixed HeLa cells using AZI2 antibody (68326-1-Ig, Clone: 2F1G5) at dilution of 1:1000 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).