

À des fins de recherche uniquement

Anticorps Monoclonal anti-THYN1

Numéro de catalogue: 68337-1-Ig



Informations de base

Numéro de catalogue: 68337-1-Ig	Numéro d'acquisition GenBank: BC006978	Méthode de purification: Purification par protéine G
Taille: 150ul , Concentration: 750 µg/ml by Nanodrop;	Identification du gène (NCBI): 29087	CloneNo.: 1E12G3
Hôte: Mouse	Nom complet: thymocyte nuclear protein 1	Dilutions recommandées: WB 1:5000-1:50000
Isotype: IgG1	MW calculé 225aa,26 kDa; 166aa,19 kDa	
Immunogen Catalog Number: AG8617	MW observés: 28 kDa	

Applications

Applications testées:
WB, ELISA

Spécificité de l'espèce:
Humain

Contrôles positifs:

WB : cellules LNCaP, cellules HeLa, cellules HepG2, cellules Jurkat, cellules K-562, cellules MOLT-4, cellules U2OS

Informations générales

THYN1, also named as THY28, is a 28-32 kDa protein which located mainly in the nucleus. It appears to correlate with induction of apoptosis. (PMID:14601557)

Stockage

Stockage:
Stocker à -20°C. Stable pendant un an après l'expédition.

Tampon de stockage:
PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3
L'aliquotage n'est pas nécessaire pour le stockage à -20C

***** Les 20ul contiennent 0,1% de BSA.**


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



Various lysates were subjected to SDS PAGE followed by western blot with 68337-1-Ig (THYN1 antibody) at dilution of 1:15000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Beta Actin Monoclonal antibody (HRP-66009) as loading control.



Various lysates were subjected to SDS PAGE followed by western blot with 68337-1-Ig (THYN1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Beta Actin Monoclonal antibody (HRP-66009) as loading control.