

À des fins de recherche uniquement

Anticorps Monoclonal anti-MED6

Numéro de catalogue: 68352-1-Ig



Informations de base

Numéro de catalogue: 68352-1-Ig	Numéro d'acquisition GenBank: BC004106	Méthode de purification: Purification par protéine A
Taille: 150ul , Concentration: 1000 µg/ml by Nanodrop;	Identification du gène (NCBI): 10001	CloneNo.: 3E4G10
Hôte: Mouse	Nom complet: mediator complex subunit 6	Dilutions recommandées: WB 1:5000-1:50000
Isotype: IgG2b	MW calculé 28 kDa	
Immunogen Catalog Number: AG7645	MW observés: 33 kDa	

Applications

Applications testées:
WB, ELISA

Spécificité de l'espèce:
Humain, rat, souris

Contrôles positifs:

WB : cellules U2OS, cellules 4T1, cellules HEK-293,
cellules HeLa, cellules HSC-T6, cellules Jurkat,
cellules LNCaP, cellules T-47D

Informations générales

MED6 is a part of mediator complex. Mediator functions as a bridge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription machinery. Mediator is recruited to promoters by direct interactions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex with RNA polymerase II and the general transcription factors.

Stockage

Stockage:
Stocker à -20°C. Stable pendant un an après l'expédition.
Tampon de stockage:
PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3
L'aliquotage n'est pas nécessaire pour le stockage à -20C

***** Les 20ul contiennent 0,1% de BSA.**


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



Various lysates were subjected to SDS PAGE followed by western blot with 68352-1-Ig (MED6 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Beta Actin Monoclonal antibody (HRP-66009) as loading control.