

À des fins de recherche uniquement

Anticorps Monoclonal anti-ANKRD54

Numéro de catalogue: 68388-1-Ig



Informations de base

Numéro de catalogue:	68388-1-Ig	Numéro d'acquisition GenBank:	BC066909	Méthode de purification:	Purification par protéine G
Taille:	150ul , Concentration: 1000 µg/ml by Nanodrop;	Identification du gène (NCBI):	129138	CloneNo.:	1E9A9
Hôte:	Mouse	Nom complet:	ankyrin repeat domain 54	Dilutions recommandées:	WB 1:2000-1:10000
Isotype:	IgG1	MW calculé	300 aa, 33 kDa		
Immunogen Catalog Number:	AG22772	MW observés:	33-35 kDa		

Applications

Applications testées:	WB, ELISA	Contrôles positifs:
Spécificité de l'espèce:	Humain, rat, souris	WB : cellules HeLa, cellules 4T1, cellules HepG2, cellules HSC-T6, cellules Jurkat, cellules K-562, cellules NIH/3T3, cellules U2OS

Informations générales

Ankyrin repeat domain 54 (ANKRD54)/Liar is a nuclear-cytoplasmic shuttling adaptor that interacts with Lyn, Btk, Tsk, and HCLS1. Activation of PKC kinases promoted nuclear export and phosphorylation of Ankrd54, and increased interaction and cytoplasmic co-localization with Lyn.

Stockage


Stockage:
Stocker à -20°C. Stable pendant un an après l'expédition.
Tampon de stockage:
PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3
L'aliquotage n'est pas nécessaire pour le stockage à -20°C

*** Les 20ul contiennent 0,1% de BSA.

For technical support and original validation data for this product please contact:
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



Various lysates were subjected to SDS PAGE followed by western blot with 68388-1-Ig (ANKRD54 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Beta Actin Monoclonal antibody (HRP-66009) as loading control.

U2OS cells were subjected to SDS PAGE followed by western blot with 68388-1-Ig (ANKRD54 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.