

À des fins de recherche uniquement

Anticorps Monoclonal anti-TBC1D23

Numéro de catalogue: 68409-1-Ig



Informations de base

| | | |
|--|---|---|
| Numéro de catalogue: 68409-1-Ig | Numéro d'acquisition GenBank: BC010221 | Méthode de purification: Purification par protéine A |
| Taille: 150ul , Concentration: 1000 µg/ml by Nanodrop; | Identification du gène (NCBI): 55773 | CloneNo.: 2A12A7 |
| Hôte: Mouse | Nom complet: TBC1 domain family, member 23 | Dilutions recommandées: WB 1:5000-1:50000 |
| Isotype: IgG2a | MW calculé: 699 aa, 78 kDa | |
| Immunogen Catalog Number: AG10727 | MW observés: 78 kDa | |

Applications

Applications testées:
WB, ELISA

Spécificité de l'espèce:
Humain, porc, rat, souris

Contrôles positifs:

WB : cellules U2OS, cellules 4T1, cellules HEK-293, cellules HSC-T6, cellules Jurkat, cellules K-562, cellules LNCaP, cellules NIH/3T3, tissu cérébral de porc

Informations générales

TBC1D23, also named as NS4ATP1, is a gene down-regulated by Pneumocystis infection. Pneumocystis pneumonia is a common opportunistic disease in AIDS patients. (PMID: 20377877) Recently it has been found that TBC1D23 gene is the most frequently mutated in MSI-H cell lines and protein expressions of TBC1D23 in tumors with mutation were down regulated.(PMID: 20824714)

Stockage

Stockage:
Stocker à -20°C. Stable pendant un an après l'expédition.

Tampon de stockage:
PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3

L'aliquotage n'est pas nécessaire pour le stockage à -20C

*** Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



Various lysates were subjected to SDS PAGE followed by western blot with 68409-1-Ig (TBC1D23 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Alpha Tubulin Monoclonal antibody (HRP-66031) as loading control.