

À des fins de recherche uniquement

Anticorps Monoclonal anti-HSDL2

Numéro de catalogue: 68414-1-Ig



Informations de base

Numéro de catalogue: 68414-1-Ig	Numéro d'acquisition GenBank: BC004331	Méthode de purification: Purification par protéine A
Taille: 150ul , Concentration: 1000 µg/ml by Nanodrop;	Identification du gène (NCBI): 84263	CloneNo.: 3D11G1
Hôte: Mouse	Nom complet: hydroxysteroid dehydrogenase like 2	Dilutions recommandées: WB 1:5000-1:50000
Isotype: IgG2b	MW calculé 45 kDa	
Immunogen Catalog Number: AG8314	MW observés: 45 kDa	

Applications

Applications testées:
WB, ELISA

Spécificité de l'espèce:
Humain, Lapin, porc

Contrôles positifs:

WB : cellules HepG2, cellules HEK-293, cellules HeLa, cellules Jurkat, cellules LNCaP, tissu cardiaque de lapin, tissu cardiaque de porc

Informations générales

Human hydroxysteroid dehydrogenase-like 2 (HSDL2) is a characterized SDR gene that not only catalyses the oxidation and reduction of multiple substrates but also regulates different metabolic and signalling pathways. Accumulating evidences suggest that HSDL2 play an important role in cancer progression. HSDL2 has promoting effects on tumor progression in papillary thyroid cancer, bladder cancer, ovarian cancer, and glioma, but has suppressing effects in cholangiocarcinoma. (PMID: 31372054, PMID: 32211805)

Stockage

Stockage:

Stocker à -20°C. Stable pendant un an après l'expédition.

Tampon de stockage:

PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3

L'aliquotage n'est pas nécessaire pour le stockage à -20C

*** Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)

E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



Various lysates were subjected to SDS PAGE followed by western blot with 68414-1-Ig (HSDL2 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.