

À des fins de recherche uniquement

Anticorps Monoclonal anti-TEAD3

Numéro de catalogue: 68464-1-Ig



Informations de base

Numéro de catalogue: 68464-1-Ig	Numéro d'acquisition GenBank: BC027877	Méthode de purification: Purification par protéine G
Taille: 150ul , Concentration: 1000 µg/ml by Nanodrop;	Identification du gène (NCBI): 7005	CloneNo.: 2E6A12
Hôte: Mouse	Nom complet: TEA domain family member 3	Dilutions recommandées: WB 1:2000-1:10000
Isotype: IgG1	MW calculé 435 aa, 49 kDa	
Immunogen Catalog Number: AG4407	MW observés: 50 kDa	

Applications

Applications testées:
WB, ELISA
Spécificité de l'espèce:
Humain, porc

Contrôles positifs:

WB : cellules HCT 116, cellules HepG2, cellules JAR,
cellules U2OS, tissu cardiaque de porc

Informations générales

TEAD3, others named transcriptional enhancer factor 5 (TEF5), a member of family of transcription factors that plays a key role in the Hippo signaling pathway, suggesting involve in organ and tumor suppression by restricting proliferation and promoting. TEAD3 contains the TEA/ATTS DNA-binding domain. Human TEAD3 has a strong mRNA expression in placenta and choriocarcinoma cells (JEG-3), but low in HeLa, HepG2 and MCF-7.

Stockage

Stockage:
Stocker à -20°C. Stable pendant un an après l'expédition.
Tampon de stockage:
PBS avec azote de sodium à 0,02 % et glycérol à 50 % pH 7,3
L'aliquotage n'est pas nécessaire pour le stockage à -20C

*** Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées



Various lysates were subjected to SDS PAGE followed by western blot with 68464-1-Ig (TEAD3 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 68464-1-Ig (TEAD3 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.