

À des fins de recherche uniquement

Anticorps Monoclonal anti-NeutraKine® IL-10



Numéro de catalogue: 69018-1-Ig **2 Publications**

Informations de base

Numéro de catalogue: 69018-1-Ig	Numéro d'acquisition GenBank: 3586	Méthode de purification: Purification par protéine G
Taille: 100ug	Identification du gène (NCBI): interleukin 10	CloneNo.: 1E4F5
Hôte: Mouse	Nom complet: interleukin 10	Dilutions recommandées: IHC 1:50-1:500
Isotype: IgG1		
Immunogen Catalog Number: HZ-1145		

Applications

Applications testées:
IHC, Neutralization, ELISA

Demandes citées:
IHC

Spécificité de l'espèce:
Humain

Espèces citées:
souris

Remarque-IHC: il est suggéré de démasquer l'antigène avec un tampon de TE buffer pH 9.0; (*) À défaut, le démasquage de l'antigène peut être effectué avec un tampon citrate pH 6,0.

Contrôles positifs:

IHC : tissu d'amygdalite humaine, tissu de cancer du poumon humain

Informations générales

Interleukin (IL)-10 is an anti-inflammatory cytokine, produced by T helper (Th) cells, macrophages, monocytes, and B cells, that plays a crucial role in preventing inflammatory and autoimmune pathologies. It downregulates the expression of Th1 cytokines, MHC class II antigens, and co-stimulatory molecules on macrophages. It also enhances B cell survival, proliferation, and antibody production. IL-10 can block NF-κB activity, and is involved in the regulation of the JAK-STAT signaling pathway. IL-10, along with its receptors, describes an important role in pathogenesis of various diseases, including infectious, inflammatory, autoimmune diseases. IL-10 mutations are associated with an increased susceptibility to HIV-1 infection and rheumatoid arthritis.

This antibody can be used to neutralize the bioactivity of IL-10.

Publications notables

Autrice	Pubmed ID	Journal	Application
Balun Li	34722505	Front Cell Dev Biol	IHC
Rui Bai	34131401	Int J Biol Sci	IHC

Stockage

Stockage:

Les anticorps lyophilisés sont stables pendant 1 an à compter de la date de réception s'ils sont stockés à une température comprise entre -20 °C et -80 °C. Après reconstitution, nous recommandons que la solution soit stockée à (4 °C) pour une courte durée ou à une température comprise entre (-20 °C) et (-80 °C) pour une longue durée. Il convient d'éviter les cycles répétés de congélation-décongélation avec les produits reconstitués.

Tampon de stockage:

PBS stérile.

L'aliquotage n'est pas nécessaire pour le stockage à -20C

*** Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)


E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Données de validation sélectionnées




Recombinant human IL-10 (Cat.NO. HZ-1145) stimulates proliferation of MC/9 cells (mouse mast cell line) in a dose-dependent manner (blue curve, refer to bottom X-left Y axis). The activity of human IL-10 (10 ng/mL) is neutralized by mouse anti-human IL-10 monoclonal antibody 69018-1-Ig at serial dose (red curve, refer to top X-right Y axis). The ND50 is typically 50-200 ng/mL. The NeutraControl mouse anti-human IL-10 monoclonal antibody 69518-1-Ig could




Recombinant human IL-10 (Cat.NO. HZ-1145) stimulates proliferation of MC/9 cells (mouse mast cell line) in a dose-dependent manner (blue curve, refer to bottom X-left Y). The activity of human IL-10 (10 ng/mL) is neutralized by mouse anti-human IL-10 monoclonal antibody 69018-1-Ig at serial dose (red curve, refer to top X-right Y). The ND50 is typically 50-200 ng/mL.

Indirect ELISA was carried out by coating recombinant Human IL-10 (Cat.NO. HZ-1145) at 70 ng/well followed by blocking and adding serial diluted IL-10 antibody 69018-1-Ig and 69518-1-Ig respectively. Signal was developed with TMB and stopped by H2SO4. Signal strength was measured by absorbance at 450 nm.



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 69018-1-Ig (NeutraKine® IL-10 antibody) at dilution of 1:200 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human tonsillitis tissue slide using 69018-1-Ig (NeutraKine® IL-10 antibody) at dilution of 1:200 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).