

À des fins de recherche uniquement

# Anticorps Monoclonal anti-NeutraControl TNF Alpha

Numéro de catalogue: 69502-1-Ig



## Informations de base

Numéro de catalogue:	69502-1-Ig	Numéro d'acquisition GenBank:	
Taille:	100ug	Identification du gène (NCBI):	7124
Hôte:	Mouse	Nom complet:	tumor necrosis factor (TNF superfamily, member 2)
Isotype:	IgG1		
Immunogen Catalog Number:	HZ-1014	Méthode de purification:	Purification par protéine G
		CloneNo.:	1A3A3

## Applications

Applications testées:  
Non-Neutralization, ELISA

Spécificité de l'espèce:  
Humain

## Informations générales

TNF, also known as TNF-alpha, or cachectin, is a multifunctional proinflammatory cytokine that belongs to the tumor necrosis factor (TNF) superfamily. It is expressed as a 26 kDa membrane bound protein and is then cleaved by TNF-alpha converting enzyme (TACE) to release the soluble 17 kDa monomer, which forms homotrimers in circulation. It is produced chiefly by activated macrophages, although it can be produced by many other cell types such as CD4+ lymphocytes, NK cells, neutrophils, mast cells, eosinophils, and neurons. It can bind to, and thus functions through its receptors TNFRSF1A/TNFR1 and TNFRSF1B/TNFB. This cytokine is involved in the regulation of a wide spectrum of biological processes including cell proliferation, differentiation, apoptosis, lipid metabolism, and coagulation. This cytokine has been implicated in a variety of diseases, including autoimmune diseases, INS resistance, and cancer.

This antibody is a neutralizing control antibody for TNF alpha, the immunogen is the same as Neutrakine 69002-1-Ig but could not neutralize human TNF alpha.

## Stockage

**Stockage:**  
Les anticorps lyophilisés sont stables pendant 1 an à compter de la date de réception s'ils sont stockés à une température comprise entre -20 °C et -80 °C. Après reconstitution, nous recommandons que la solution soit stockée à (4 °C) pour une courte durée ou à une température comprise entre (-20 °C) et (-80 °C) pour une longue durée. Il convient d'éviter les cycles répétés de congélation-décongélation avec les produits reconstitués.


**Tampon de stockage:**  
PBS stérile.  
**L'aliquotage n'est pas nécessaire pour le stockage à -20°C**

\*\*\* Les 20ul contiennent 0,1% de BSA.

For technical support and original validation data for this product please contact:  
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free  
in USA), or 1(312) 455-8498 (outside USA)      E: proteintech@ptglab.com  
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Données de validation sélectionnées



Recombinant human TNF alpha (Cat.NO. HZ-1014) inhibits the growth of L-929 cell line in the presence of actinomycin D in a dose dependent manner (blue curve, refer to bottom X-left Y). The activity elicited by recombinant human TNF alpha (0.05 ng/mL HZ-1014) is neutralized by mouse anti-human TNF alpha monoclonal antibody 69002-1-Ig at serial dose (red curve, refer to top X-right Y). The ND<sub>50</sub> is typically 10-50 ng/mL. The NeutraControl mouse anti-human TNF alpha

Indirect ELISA was carried out by coating recombinant Human TNF Alpha (Cat.NO. HZ-1014) at 70 ng/well followed by blocking and adding serial diluted TNF Alpha antibody 69002-1-Ig and 69502-1-Ig respectively. Signal was developed with TMB and stopped by H<sub>2</sub>SO<sub>4</sub>. Signal strength was measured by absorbance at 450 nm.