

À des fins de recherche uniquement

# Anticorps Monoclonal anti-NeutraControl TGF beta 1



Numéro de catalogue: 69512-1-Ig

## Informations de base

<b>Numéro de catalogue:</b> 69512-1-Ig	<b>Numéro d'acquisition GenBank:</b> 7040	<b>Méthode de purification:</b> Purification par protéine G
<b>Taille:</b> 100ug, Concentration: 1000 µg/ml by Nanodrop;	<b>Identification du gène (NCBI):</b> 7040	<b>CloneNo.:</b> 1D3D11
<b>Hôte:</b> Mouse	<b>Nom complet:</b> transforming growth factor, beta 1	
<b>Isotype:</b> IgG1		
<b>Immunogen Catalog Number:</b> HZ-1011		

## Applications

**Applications testées:**  
Non-Neutralization, ELISA

**Spécificité de l'espèce:**  
Humain

## Informations générales

TGFB, also named as LAP and TGFB1, is a multifunctional peptide that controls proliferation, differentiation, and other functions in many cell types. TGFB acts synergistically with TGFA in inducing transformation. It also acts as a negative autocrine growth factor. Dysregulation of TGFB activation and signaling may result in apoptosis. Many cells synthesize TGFB and almost all of them have specific receptors for it. TGFB positively and negatively regulates many other growth factors. It plays an important role in bone remodeling as it is a potent stimulator of osteoblastic bone formation, causing chemotaxis, proliferation and differentiation in committed osteoblasts. It is highly expressed in bone. Mutation of TGFB are the cause of Camurati-Engelmann disease (CED) which known as progressive diaphyseal dysplasia 1 (DPD1).

This antibody is a neutralizing control antibody for TGF beta 1, the immunogen is the same as Neutrakine 69012-1-Ig but could not neutralize human TGF beta 1.

## Stockage

**Stockage:**  
Les anticorps lyophilisés sont stables pendant 1 an à compter de la date de réception s'ils sont stockés à une température comprise entre -20 °C et -80 °C. Après reconstitution, nous recommandons que la solution soit stockée à (4 °C) pour une courte durée ou à une température comprise entre (-20 °C) et (-80 °C) pour une longue durée. Il convient d'éviter les cycles répétés de congélation-décongélation avec les produits reconstitués.

**Tampon de stockage:**  
PBS stérile.

L'aliquotage n'est pas nécessaire pour le stockage à -20C

\*\*\* Les 20ul contiennent 0,1% de BSA.



For technical support and original validation data for this product please contact:

T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free in USA), or 1(312) 455-8498 (outside USA)

E: [proteintech@ptglab.com](mailto:proteintech@ptglab.com)  
W: [ptglab.com](http://ptglab.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Données de validation sélectionnées



Recombinant human TGF beta 1 (Cat.NO. HZ-1011) inhibits IL-4 induced proliferation of mouse HT-2 cells in a dose dependent manner (blue curve, refer to bottom X-left Y). The activity of human TGF beta 1 (0.5 ng/mL HZ-1011) is neutralized by mouse anti-human TGF beta 1 antibody 69012-1-Ig at serial dose (red curve, refer to top X-right Y). The ND50 is typically 1-5 µg/mL. The NeutraControl mouse anti-human TGF beta 1 monoclonal antibody 69512-1-Ig could recognize human TGF beta 1,

Indirect ELISA was carried out by coating recombinant Human TGF beta 1 (Cat.NO. HZ-1011) at 70 ng/well followed by blocking and adding serial diluted TGF beta 1 antibody 69012-1-Ig and 69512-1-Ig respectively. Signal was developed with TMB and stopped by H<sub>2</sub>SO<sub>4</sub>. Signal strength was measured by absorbance at 450 nm.