Anticorps Recombinant de lapin antiproteintech SARS-CoV-2 Nucleocapsid Phosphoprotein



Informations de base	Numéro de catalogue: 80026-1-RR	Numéro d'acquisition GenBank: NC_045512	Méthode de purification: Purification par protéine A
	Taille: 100ul , Concentration: 1000 µg/ml by Nanodrop; Hôte: Lapin	Identification du gène (NCBI): 43740575	CloneNo.: 4B9
		Nom complet: COVID-19 N Protein	Dilutions recommandées: WB 1:5000-1:50000
	lsotype: IgG		
	Immunogen Catalog Number: AG30676		
Applications	Applications testées: WB,ELISA	Contrôles positifs: WB : Phosphoprotéine nucléocapside eucaryote,	
	Spécificité de l'espèce: Virus		
Informations générales	The nucleocapsid (N) protein has multiple functions including formation of nucleocapsids, signal transduction virus budding, RNA replication, and mRNA transcription. N protein is an important antigen for coronavirus, and it is normally highly conserved, with a molecular weight of about 50 kDa. it can be used as a marker in diagnostic assays due to its high immunogenicity (PMID: 32416961, PMID: 32235387). A sandwich ELISA for COVID-19 N Protein can be assembled by using 80027-1-RR as capture antibody and conjugated 80026-1-RR for detection.		
Stockage	Stockage: Stocker à -20 °C. Tampon de stockage: PBS avec azoture de sodium à 0,02 %		
*** Les 20ul contiennent 0,1% de BSA	L'aliquotage n'est pas nécessaire pou	r le stockage à -20C	


For technical support and original validation data for this product please contact: T: 1 (888) 4PTGLAB (1-888-478-4522) (toll free E: proteintech@ptglab.com in USA), or 1(312) 455-8498 (outside USA) W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Antibodies | ELISA kits | Proteins

www.ptglab.com


Données de validation sélectionnées




E.coli expressed SARS-CoV-2 Nucleocapsid Phosphoprotein (Cat.NO. Ag30676) was subjected to SDS-PAGE followed by western blot with 80027-1-RR and 80026-1-RR at various work concentration.



Eukaryotic expressed SARS-CoV-2 Nucleocapsid Phosphoprotein was subjected to SDS-PAGE followed by western blot with 80027-1-RR and 80026-1-RR at various work concentration.



Indirect ELISA was carried out by coating eukaryotic expressed N protein at 70 ng/well followed by blocking and adding serial diluted primary antibody 80026-1-RR and 80027-1-RR respectively. Signal was developed with TMB and stopped by H2SO4. Signal strength was measured by absorbance at 450 nm.



Sandwich ELISA was carried out by coating 80027-1-RR at 80 ng/well followed by blocking and adding different concentration of eukaryotic expressed N protein (0.5-1000 ng/mL). HRPconjugated80026-1-RR was used at 1 µg/mL for detection. Signal was developed with TMB and stopped by H2SO4. Signal strength was measured by absorbance at 450 nm.