

À des fins de recherche uniquement

Anticorps Recombinant de lapin anti-CD31



Numéro de catalogue: 80530-1-RR

Informations de base

Numéro de catalogue:	BC022512	Méthode de purification:
Taille:	Identification du gène (NCBI):	Purification par protéine A
100ul , Concentration: 1000 µg/ml by Nanodrop;	5175	CloneNo.:
Hôte:	Nom complet:	4N13
Lapin	platelet/endothelial cell adhesion molecule	Dilutions recommandées:
Isotype:	MW calculé:	WB 1:5000-1:50000
IgG	83 kDa	IF 1:250-1:1000
Immunogen Catalog Number:	MW observés:	
AG1787	130 kDa	

Applications

Applications testées:	Contrôles positifs:
IF, WB, ELISA	WB : plaquettes de sang périphérique humain, cellules Jurkat, cellules MOLT-4
Spécificité de l'espèce:	
Humain, souris	IF : cellules bEnd.3,

Informations générales

Platelet endothelial cell adhesion molecule-1 (PECAM-1, CD31) is a member of the immunoglobulin gene superfamily of cell adhesion molecules. CD31 is a transmembrane glycoprotein that is highly expressed on the surface of the endothelium, making up a large portion of its intracellular junctions. PECAM-1 is also present on the surface of hematopoietic cells and immune cells including platelets, monocytes, neutrophils, natural killer cells, megakaryocytes and some types of T-cell (PMID: 9011572). As well as its role in cell-cell adhesion, PECAM-1 functions as a signaling receptor, and is involved in important physiological events such as nitric oxide production, regulation of T-cell immunity and tolerance, leukocyte transendothelial migration and inflammation and angiogenesis (PMID: 21183735; 20978210; 17872453; 20634489).

Stockage


Stockage:
Stocker à -20 °C.
Tampon de stockage:
PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3
L'aliquotage n'est pas nécessaire pour le stockage à -20°C

*** Les 20ul contiennent 0,1% de BSA.


For technical support and original validation data for this product please contact:
T: 1(888) 4PTGLAB (1-888-478-4522) (toll free
in USA), or 1(312) 455-8498 (outside USA) E: proteintech@ptglab.com
W: ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Données de validation sélectionnées



Untreated and PNGase F-treated lysates of human peripheral blood platelets were subjected to SDS PAGE followed by western blot with 80530-1-RR (CD31 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 80530-1-RR (CD31 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed bEnd.3 cells using CD31 antibody (80530-1-RR, Clone: 4N13) at dilution of 1:500 and Coralite® 488-Conjugated AffiniPure Goat Anti-Rabbit IgG(H+L), Alpha Tubulin antibody (66031-1 Ig, Clone: 1E4C11, red).