

# Anticorps Recombinant de lapin anti-YAP1

Numéro de catalogue: 81090-1-RR

## Informations de base

Numéro de catalogue:	Numéro d'acquisition GenBank:	Méthode de purification:
81090-1-RR	BC038235	Purification par protéine A
Taille:	Identification du gène (NCBI):	CloneNo.:
100ul , Concentration: 1000 µg/ml by Nanodrop;	10413	5E3
Hôte:	Nom complet:	Dilutions recommandées:
Lapin	Yes-associated protein 1, 65kDa	WB 1:5000-1:50000
Isotype:	MW calculé	IHC 1:250-1:1000
IgG	504 aa, 54 kDa	
Immunogen Catalog Number:	MW observés:	
AG4510	65-70 kDa	

## Applications

Applications testées:	Contrôles positifs:
IHC, WB, ELISA	WB: cellules HeLa,
Spécificité de l'espèce:	IHC : tissu de cancer du côlon humain, tissu de muscle squelettique de souris
Humain, rat, souris	

**Remarque-IHC:** il est suggéré de démasquer l'antigène avec un tampon de TE buffer pH 9,0; (\*) À défaut, le démasquage de l'antigène peut être effectué avec un tampon citrate pH 6,0.

## Informations générales


Yes-associated protein 1 (YAP1) is a transcriptional regulator which can act both as a coactivator and a corepressor and is the critical downstream regulatory target in the Hippo signaling pathway that plays a pivotal role in organ size control and tumor suppression by restricting proliferation and promoting apoptosis. The core of this pathway is composed of a kinase cascade wherein STK3/MST2 and STK4/MST1, in complex with its regulatory protein SAV1, phosphorylates and activates LATS1/2 in complex with its regulatory protein MOB1, which in turn phosphorylates and inactivates YAP1 oncprotein and WWTR1/TAZ. Plays a key role to control cell proliferation in response to cell contact. Phosphorylation of YAP1 by LATS1/2 inhibits its translocation into the nucleus to regulate cellular genes important for cell proliferation, cell death, and cell migration. The presence of TEAD transcription factors are required for it to stimulate gene expression, cell growth, anchorage-independent growth, and epithelial-mesenchymal transition (EMT) induction. Isoform 2 and isoform 3 can activate the C-terminal fragment (CTF) of ERBB4 (isoform 3). Increased expression seen in some liver and prostate cancers. Isoforms lacking the transactivation domain found in striatal neurons of patients with Huntington disease (at protein level). It is activated by phosphorylation and degraded by ubiquitination (20048001). The calculated molecular weight of YAP1 is 54 kDa, but phosphorylated YAP1 is about 65-70 kDa. (PMID: 26695440)

## Stockage




Stockage:  
Stocker à -20 °C.  
Tampon de stockage:  
PBS avec azoture de sodium à 0,02 % et glycérol à 50 % pH 7,3  
L'aliquotage n'est pas nécessaire pour le stockage à -20°C

\*\*\* Les 20ul contiennent 0,1% de BSA.


## Données de validation sélectionnées



Various lysates were subjected to SDS PAGE followed by western blot with 81090-1-RR (YAP1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded mouse skeletal muscle tissue slide using 81090-1-RR (YAP1 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human colon cancer tissue slide using 81090-1-RR (YAP1 antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).